

Quantum Continuum Mechanics Made Simple

Tim Gould,¹ Georg Jansen,² I. V. Tokatly,^{3,4} and John F. Dobson¹

¹*Qld Micro- and Nanotechnology Centre, Griffith University, Nathan, Qld 4111, Australia*

²*Fakultät für Chemie, Universität Duisburg-Essen, 45117 Essen, Germany*

³*ETSF Scientific Development Centre, Departamento de Física de Materiales, Universidad del País Vasco UPV/EHU, Av. Tolosa 72, E-20018 San Sebastián, Spain*

⁴*IKERBASQUE, Basque Foundation for Science, E-48011, Bilbao, Spain*

In this paper we further explore and develop the quantum continuum mechanics (CM) of [Tao *et al*, PRL**103**,086401] with the aim of making it simpler to use in practice. Our simplifications relate to the non-interacting part of the CM equations, and primarily refer to practical implementations in which the groundstate stress tensor is approximated by its Kohn-Sham version. We use the simplified approach to directly prove the exactness of CM for one-electron systems via an orthonormal formulation. This proof sheds light on certain physical considerations contained in the CM theory and their implication on CM-based approximations. The one-electron proof then motivates an approximation to the CM (exact under certain conditions) expanded on the wavefunctions of the Kohn-Sham (KS) equations. Particular attention is paid to the relationships between transitions from occupied to unoccupied KS orbitals and their approximations under the CM. We also demonstrate the simplified CM semi-analytically on an example system.

PACS numbers: 31.15.E-,31.15.ee,31.15.xg,31.15.ap

I. INTRODUCTION

The study of the continuum mechanics (fluid dynamics) of quantum electron fluids is almost as old as quantum mechanics itself. The Thomas-Fermi model and Madelung dynamics¹ are two very early examples. Unfortunately standard approaches to Continuum Mechanics become too inaccurate or too complex when applied to larger, real systems such as molecules and atoms, although a recent development² makes atomic systems more tractable. Recent work³⁻⁷ on Continuum Mechanics (CM) in a moving Lagrangian frame has led to the development of a sophisticated approach for describing linear perturbations to many-electron systems from groundstate properties only. This approach appears able to bridge the gap between speed and accuracy required by modern *ab initio* calculations.

The CM provides an efficient⁷ alternative to full time-dependent density functional theory (tdDFT)⁸⁻¹³ calculations. In its general form it can, in principle, be used to evaluate the transition frequencies and currents of a real many-electron quantum system using the one- and two-particle density matrices. In the formalism presented here we restrict to a more limited form that takes, as input, groundstate properties obtained from a Kohn-Sham⁸ calculation, and approximates small changes to the groundstate via a continuum approach. This restriction to groundstate KS properties only comes at the expense of having to deal with higher order mixed derivatives (up to four derivatives with three indices). Initial indications suggest, however, that it is both tractable and valid in both model systems,^{5,6} as well as the difficult and geometrically very different case of two interacting two-dimensional electron gas layers.⁷

Like Madelung dynamics, the recent CM approach describes the behavior of the fluid displacement vector $\mathbf{u}$ from which the current and changes to the density can be described. As the independent-electron density response $\hat{\chi}_0$ of a system can be obtained from $\mathbf{u}$, a “CM-dRPA” correlation energy functional⁷ has been developed that uses the direct Random Phase Approximation (dRPA) but bypasses the need for unoccupied orbitals by working in the CM directly via $\mathbf{u}$. The CM-dRPA can be considered a “third-rung” functional according to the “Jacob’s Ladder” classification scheme of Perdew *et al.*¹⁴ The functional involves calculating the bare response via the CM scheme and solving for the interacting response $\hat{\chi}_\lambda$ under the dRPA where additional interactions are treated at time-dependent Hartree level via $\hat{\chi}_\lambda = \hat{\chi}_0 + \lambda \hat{\chi}_0 \hat{v} \hat{\chi}_\lambda$. The response functions $\hat{\chi}_0$ and $\hat{\chi}_\lambda$ can then be used to calculate

correlation energies. Here λ is the strength of the interactions and must be integrated over to obtain the kinetic contribution to the correlation energy.

One particular area where it is hoped that the CM will prove broadly useful is in the evaluation of van der Waals physics, where long-range correlation is important. Here local-density based techniques^{15–17} and even vdW-adapted approximations like the vdW-DF^{18–21} run into difficulties (see ref. 7 and ref. 22 for discussion) because of their use of local or pairwise approximations.

The strong theoretical relationships between the CM and the KS-like system it approximates, arising from its derivation from a formal, moving Lagrange frame,⁴ is somewhat hidden in the current prescription, especially to those used to working with orbitals and the Schrödinger equation. While relationships can be established (such as sum rules) between $\mathbf{u}$ and common groundstate properties of interest, they do not always come naturally from the original formulation.

Via adaptation and exploration of its theoretical form the CM offers a wider scope for further investigation, as follows:

Firstly, in their original form⁵ the CM equations were complicated. In the first application to vdW physics,⁷ a more compact, simpler and more symmetric but equivalent form was used (equations 3-5 of ref. 7). The derivation of this simplified form was not given in ref. 7, and is presented here for the first time along with a discussion of the particular version of the groundstate stress tensor that makes this simplified version of the CM theory possible.

Secondly, the standard displacement vector $\mathbf{u}$ has some undesirable properties in Coulomb-localised systems such as atoms and molecules, including divergence in the tail of the density distribution. To deal with this we reformulate the CM in terms of an “orthonormal displacement vector” $\boldsymbol{\xi} = \sqrt{n^0}\mathbf{u}$ (touched on in equation (68) of ref. 6) where n^0 is the electron density of the groundstate. This alternate approach opens the method up to expansion in basis sets with decaying tails, such as the widely used²³ Gaussian-Type Orbitals (GTOs) and Slater-Type Orbitals (STOs) which are only valid in the expansion of decaying functions.

Thirdly, this reformulated CM can be used to demonstrate, directly from the Schrödinger equation, that the CM gives the exact bare, linear response $\hat{\chi}_0$ for one-electron systems. While this relationship is established and demonstrated in earlier works (see equations (41)-(46) in ref. 4, equation (98) and Appendix D in ref. 5) a proof direct from the Schrödinger equation has not so far appeared. This relationship is very important, having as a con-

sequence that properties dependent on $\hat{\chi}_0$ such as the (dRPA) asymptotic van der Waals interaction between atoms, are exactly reproduced by CM-dRPA theory for one-electron systems (and two-electron systems with equal groundstate densities of spin up and down electrons). The direct proof demonstrates how the CM relates exactly to the first-order change of the one-electron orbital/wavefunction.

Fourthly, the one-electron case motivates a second, approximate reformulation of the tensor CM into a scalar system. Here the displacement $\mathbf{u}$ is approximated as the gradient of a scalar function s . This approach simplifies the CM equations at the expense of accuracy in general systems but is exact for one-electron and one-dimensional systems. By then expanding $\sqrt{n^0}s$ on the set of KS orbitals we also uncover some of the physics of the CM in the asymptotic tail regions of the density, where the electrons behave like a one-electron system.

Finally we illustrate the work on an example system: a one-dimensional, non-interacting Harmonic potential model. Here the one-electron case can be solved analytically while many of the terms in the many-electron case can be solved for analytically or using near exact quadrature, minimising numerical error. The scalar approximation is an exact reformulation of the CM in this example.

A. Notation

In this paper we work entirely in atomic coordinates where $m_{e^-} = \hbar = e^2/(4\pi\epsilon_0) = 1$ such that energies are in Hartree and distances in Bohr radii. We treat energies as frequencies with the division by $\hbar$ implicit.

Greek sub/super-scripts are used to refer to Cartesian (x , y and z) coordinates and are summed over if repeated. We use the derivative operator notation $[\partial_\mu f(\mathbf{r})] \equiv [\partial f(\mathbf{r})/\partial r_\mu]$.

Cartesian tensors are written with sans serif letters (eg. $\mathbb{T}$), while Cartesian vectors appear in bold (eg. $\mathbf{v}$). Their elements are typically given as $T_{\mu\nu}$ and v_α respectively. More general matrices use double-line Roman letters (eg. $\mathbb{M}$) and should be considered square unless otherwise noted.

Operators involve a hat eg. $\hat{O}$. If a derivative appears in an operator it can be considered to act *entirely* to the right unless surrounded by *square* brackets, but will act through other brackets. Thus $[\partial_\alpha v_\alpha] \equiv [\nabla \cdot \mathbf{v}]$, $\partial_\mu f \equiv [\partial_\mu f] + f\partial_\mu$ and $(\partial_\mu f)g \equiv [\partial_\mu f]g + f[\partial_\mu g] + fg\partial_\mu$.

Comma-led subscripts will sometimes be used to represent derivatives $[\partial_\mu f_\nu] \equiv f_{\nu,\mu}$.

In the context of KS orbitals j or k can typically be any orbital but we reserve i for occupied orbitals only and a for unoccupied orbitals only such that $\sum_i \equiv \sum_{i \text{ occ}}$ and $\sum_{ai} \equiv \sum_{i \text{ occ}} \sum_{a \text{ unocc}}$.

II. ORIGINAL VS OPERATOR FORMS OF CM

Let us define a Kohn-Sham (KS) system with a KS potential V^{KS} (approximate or otherwise). The one-electron Hamiltonian is thus

$$\left\{ -\frac{1}{2}\nabla^2 + V^{\text{KS}}(\mathbf{r}) \right\} \psi_j(\mathbf{r}) = \epsilon_j \psi_j(\mathbf{r}) \quad (1)$$

where $\psi_j(\mathbf{r}) = \langle \mathbf{r} | j \rangle$ is a one-electron orbital which are orthonormal under $\int d\mathbf{r} \psi_j^*(\mathbf{r}) \psi_k(\mathbf{r}) \equiv \langle j | k \rangle = \delta_{jk}$ and where ϵ_j is its Kohn-Sham eigenvalue.

The groundstate density n^0 is, in general, defined via $n(\mathbf{r}) = \rho(\mathbf{r}, \mathbf{r})$ where the one-electron density matrix $\rho(\mathbf{r}, \mathbf{r}')$ is discussed later. In a KS system the density becomes

$$n^0(\mathbf{r}) = \sum_j f_j \psi_j^*(\mathbf{r}) \psi_j(\mathbf{r}) \quad (2)$$

where f_j is the occupation of orbital $|j\rangle$ defined as 1 for orbitals with $\epsilon_j < \epsilon_F$ and 0 otherwise where ϵ_F is the Fermi energy. ϵ_F should be chosen to ensure that $\int n^0(\mathbf{r}) d\mathbf{r} = N_e$ where N_e is the total number of electrons.

A. Kinetic stress tensor

For the stress tensor $\mathbb{T}$ corresponding to a one-electron density matrix $\rho(\mathbf{r}, \mathbf{r}')$ in the absence of a current we choose the following definition, which we motivate and derive in Appendix A:

$$\begin{aligned} T_{\mu\nu}(\mathbf{r}) = & \frac{1}{2} (\partial_\mu \partial'_\nu + \partial'_\mu \partial_\nu) \rho(\mathbf{r}, \mathbf{r}')|_{\mathbf{r}=\mathbf{r}'} \\ & - \frac{1}{4} \partial_\mu \partial_\nu n_0(\mathbf{r}). \end{aligned} \quad (3)$$

This particular definition seems to allow for the most compact presentation of the CM eigen-equations (18).

Inserting the density matrix from the independent-electron Kohn-Sham groundstate we obtain

$$T_{\mu\nu}^0 = \Re \sum_j f_j [\partial_\mu \psi_j^*] [\partial_\nu \psi_j] - \frac{1}{4} \partial_\mu \partial_\nu n^0 \quad (4)$$

$$\equiv \frac{1}{2} \Re \sum_j f_j ([\partial_\mu \psi_j^*] [\partial_\nu \psi_j] - \psi_j^* [\partial_\mu \partial_\nu \psi_j]) \quad (5)$$

where terms are functions of $\mathbf{r}$ only. Unless otherwise noted we subsequently restrict ourselves to this form. The kinetic stress tensor $\mathbf{T}^0$ is real symmetric [$T_{\mu\nu}^0(\mathbf{r}) = T_{\nu\mu}^0(\mathbf{r})$] and obeys the groundstate force balance condition $[\partial_\alpha T_{\alpha\mu}^0] = -n^0 [\partial_\mu V^{\text{KS}}]$.

In Appendix A we discuss the origins of the form used here and reasons for the difference between our forms of $\mathbf{T}$ and $\mathbf{T}^0$ and the form defined in equation 17 of ref. 5 [equation 13 here, leading to our equation 14 for the Kohn-Sham groundstate.]

B. Linear CM

In general, there is a linear relationship between the first-order time-dependent fluid displacement vector $\mathbf{u}(\mathbf{r}, t)$, and a small change to the external potential defined via its force density per electron $\mathbf{F}^{1\text{Ext}}(\mathbf{r}, t) \equiv n^0(\mathbf{r}) \nabla V^{1\text{Ext}}(\mathbf{r}, t)$. Here $\mathbf{u}$ is related to the electron current through²⁴

$$\mathbf{j}(\mathbf{r}, t) = \partial_t n^0(\mathbf{r}) \mathbf{u}(\mathbf{r}, t), \quad (6a)$$

$$\partial_t n^1(\mathbf{r}, t) = -\nabla \cdot \mathbf{j}(\mathbf{r}, t) \quad (6b)$$

from which it follows that a solution is

$$n^1(\mathbf{r}) = -\nabla \cdot n^0(\mathbf{r}) \mathbf{u}(\mathbf{r}). \quad (7)$$

Linear CM under a Kohn-Sham groundstate provides this relationship approximately in the following form for a time-periodic potential of frequency ω ^{5,6}

$$\omega^2 n^0 \mathbf{u} = \mathbf{F}^{1\text{Pot}} + \mathbf{F}^{1\text{Kin}} + \mathbf{F}^{1\text{Ext}} + \mathbf{F}^{1\text{Int}}, \quad (8)$$

where $\mathbf{F}^{1\text{Int}}$ includes the internal dynamic electron-electron interactions and is discussed later. The remaining terms are: $\mathbf{F}^{1\text{Kin}} = -\hat{K} \mathbf{u}$ where⁷

$$\hat{K}_{\mu\nu} = \partial_\alpha T_{\mu\nu}^0 \partial_\alpha + \partial_\alpha T_{\alpha\nu}^0 \partial_\mu + \partial_\nu T_{\alpha\mu}^0 \partial_\alpha - \frac{1}{4} \partial_\nu \partial_\alpha n^0 \partial_\alpha \partial_\mu \quad (9)$$

is an Hermitian operator, a full derivation of which appears in appendix B; And $\mathbf{F}^{1\text{Pot}} = n^0[\nabla \otimes \nabla V^{\text{KS}}] \cdot \mathbf{u}$. Putting them together in the form $\mathbf{F}^{1\text{Pot}} + \mathbf{F}^{1\text{Kin}} = \hat{\mathbf{R}}\mathbf{u}$ leads to

$$(\omega^2 n^0 - \hat{\mathbf{R}})\mathbf{u} = F^{1\text{Ext}} + F^{1\text{Int}}, \quad (10)$$

$$\hat{R}_{\mu\nu} = n^0 V_{,\mu\nu}^{\text{KS}} - \hat{K}_{\mu\nu} \quad (11)$$

where all terms but ω vary with $\mathbf{r}$. The operator $\hat{\mathbf{R}}$ is manifestly Hermitian, and can be shown to be positive definite.

The original papers^{5,6} establishing the linearised CM theory give $\mathbf{F}^{1\text{Kin}}$ [our (9)] in a less compact but exactly equivalent manner. The expression for $\mathbf{F}^{1\text{Kin}}$ appearing in equation 14 of ref. 5 (equation 53 of ref. 6) is (where we have rearranged the order)

$$\begin{aligned} -F_{\mu}^{1\text{Kin}} = & \partial_{\alpha}(2\bar{T}_{\mu\nu}^0 U_{\nu\alpha} + \bar{T}_{\nu\alpha}^0 \partial_{\mu} u_{\nu}) \\ & + \frac{1}{4} \partial_{\nu} \{ [\partial_{\nu} n^0] \partial_{\mu} + [\partial_{\mu} n^0] \partial_{\nu} - \partial_{\mu} n^0 \partial_{\nu} \} \nabla \cdot \mathbf{u} \\ & + \frac{1}{2} \partial_{\nu} \{ [\nabla^2 n^0] U_{\mu\nu} - \partial_{\mu} ([\partial_{\alpha} n^0] U_{\nu\alpha}) \}. \end{aligned} \quad (12)$$

Here $U_{\mu\nu} = \frac{1}{2}[\partial_{\nu} u_{\mu} + \partial_{\mu} u_{\nu}]$ and their kinetic stress tensor is [equation (17) in ref. 6]

$$\begin{aligned} \bar{T}_{\mu\nu}(\mathbf{r}) = & \frac{1}{2} [(\partial_{\mu} \partial'_{\nu} + \partial'_{\mu} \partial_{\nu}) \rho(\mathbf{r}, \mathbf{r}')]_{\mathbf{r}=\mathbf{r}'} \\ & - \frac{1}{4} \delta_{\mu\nu} [\nabla^2 n_0(\mathbf{r})]. \end{aligned} \quad (13)$$

For an independent-electron Kohn-Sham groundstate this takes the form

$$\bar{T}_{\mu\nu}^0 = \Re \sum_j f_j [\partial_{\mu} \psi_j^*] [\partial_{\nu} \psi_j] - \frac{1}{4} \delta_{\mu\nu} [\nabla^2 n^0] \quad (14)$$

where terms depend on $\mathbf{r}$ only. The second components in (13) and (14) differ from ours [compare (3) and (4)]. Although it is not immediately obvious, the force (12) is equivalent to an Hermitian operator acting on $\mathbf{u}$. By contrast the Hermiticity of our operator (9) is apparent on inspection.

Equation (9) defines the same force as (12) but in a much simplified manner. Much of this simplification comes from the different choice of kinetic stress tensor, given by (3) and (4) in our work and by (13) and (14) in refs 5 and 6. In fact there is some ambiguity in the definition of the quantal stress tensor. In Appendix A we use the Wigner distribution to motivate our present form, which is then used to derive (9) in Appendix B.

III. CM RESPONSE FUNCTIONS

In (8) $\mathbf{F}^{\text{1Int}}$ takes into account the small change in the Kohn-Sham potential arising from the change in density, while $\hat{\mathbf{R}}\mathbf{u}$ deals with kinetic and potential physics. The dRPA is equivalent to setting

$$\mathbf{F}^{\text{1Int}}(\mathbf{r}) = [\hat{\mathbf{Q}}\mathbf{u}](\mathbf{r}) \quad (15)$$

$$\begin{aligned} &\equiv -n^0(\mathbf{r})\nabla \int \frac{d\mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|} [\nabla' \cdot n^0(\mathbf{r}')\mathbf{u}(\mathbf{r}')] \\ &\equiv \int d\mathbf{r}' \mathbf{Q}(\mathbf{r}, \mathbf{r}') \cdot \mathbf{u}(\mathbf{r}') \end{aligned} \quad (16)$$

where

$$Q_{\mu\nu}(\mathbf{r}, \mathbf{r}') = n^0(\mathbf{r})n^0(\mathbf{r}') \left[\partial_\mu \partial'_\nu \frac{1}{|\mathbf{r} - \mathbf{r}'|} \right]. \quad (17)$$

In the absence of an external field $\mathbf{F}^{\text{1Ext}} = \mathbf{0}$ we can find eigen-mode pairs Ω_N and $\mathbf{u}_N$ (or $\Omega_{N\lambda}$ and $\mathbf{u}_{N\lambda}$) through solutions of

$$\Omega_N^2 n^0 \mathbf{u}_N = \hat{\mathbf{R}}\mathbf{u}_N, \quad (18)$$

$$\Omega_{N\lambda}^2 n^0 \mathbf{u}_{N\lambda} = \hat{\mathbf{R}}\mathbf{u}_{N\lambda} + \lambda \hat{\mathbf{Q}}\mathbf{u}_{N\lambda} \quad (19)$$

where (19) includes the internal interactions $\lambda \hat{\mathbf{Q}}$ at coupling strength λ while we use the short-hand $\Omega_N = \Omega_{N0}$ and $\mathbf{u}_N = \mathbf{u}_{N0}$ for the non-interacting case. Because $\hat{\mathbf{R}}$ and $\hat{\mathbf{Q}}$ are Hermitian and n^0 is symmetric and positive definite, eigensolutions can be found that obey the orthonormality condition

$$\int d\mathbf{r} n^0 \mathbf{u}_{N\lambda}^* \cdot \mathbf{u}_{M\lambda} = \delta_{NM}. \quad (20)$$

The set $\{\mathbf{u}_{N\lambda}\}$ is also guaranteed to be complete over a finite basis when $\hat{\mathbf{R}} + \lambda \hat{\mathbf{Q}}$ is represented in the same finite basis. Furthermore the eigen-values $\Omega_{N\lambda}$ can be shown⁶ to be positive. Typically we also sort the modes such that $\Omega_{N+1\lambda} \geq \Omega_{N\lambda}$ where $N \geq 1$ labels the CM mode with the N th lowest energy. The displacement $\mathbf{u}_{N\lambda}$ corresponds to a transition density mode (the meaning will become clearer later) defined as

$$d_{N\lambda}(\mathbf{r}) = -\nabla \cdot n^0(\mathbf{r})\mathbf{u}_{N\lambda}(\mathbf{r}). \quad (21)$$

If the external force density $\mathbf{F}^{1\text{Ext}}$ is reintroduced we can expand the solution of (8) at interaction strength λ on the basis $\{\mathbf{u}_{N\lambda}\}$ such that $\mathbf{u} = \sum_N c_{N\lambda} \mathbf{u}_{N\lambda}$. Here

$$c_{N\lambda} = \frac{\int d\mathbf{r} n^0(\mathbf{r}) \mathbf{u}_{N\lambda}^*(\mathbf{r}) \cdot \mathbf{F}^{1\text{Ext}}(\mathbf{r})}{\Omega_{N\lambda}^2 - \omega^2} \quad (22)$$

when $\mathbf{F}^{1\text{Ext}}$ is time-periodic with frequency ω . The change in density (7) can thus be expanded on (21) as

$$n_\lambda^1(\mathbf{r}, \omega) = \sum_N c_{N\lambda} d_{N\lambda}(\mathbf{r}), \quad (23)$$

where the sum is over all eigen-solutions.

The density response χ_λ of a system is defined as the change in density in response to a $\delta(\mathbf{r} - \mathbf{r}')$ potential at a frequency ω with internal interactions at strength λ . This corresponds to an external force $\mathbf{F}^{1\text{Ext}} = n^0 \nabla \delta(\mathbf{r} - \mathbf{r}') e^{-i\omega t}$ and internal force $\mathbf{F}^{1\text{Int}} = \lambda \hat{\mathbf{Q}} \mathbf{u}(\omega) e^{-i\omega t}$. Thus the response takes the form $\hat{\chi}_\lambda(t) = \hat{\chi}_\lambda(\omega) e^{-i\omega t}$ where $\chi_\lambda(\mathbf{r}, \mathbf{r}'; \omega) = -\sum_N \frac{d_{N\lambda}^*(\mathbf{r}) d_{N\lambda}(\mathbf{r}')}{\Omega_{N\lambda}^2 - \omega^2}$. Typically it is easier to work with responses at imaginary frequency $\omega = i\sigma$ such that

$$\chi_0(\mathbf{r}, \mathbf{r}'; i\sigma) = -\Re \sum_N \frac{d_N^*(\mathbf{r}) d_N(\mathbf{r}')}{\sigma^2 + \Omega_N^2}, \quad (24)$$

$$\chi_\lambda(\mathbf{r}, \mathbf{r}'; i\sigma) = -\Re \sum_N \frac{d_{N\lambda}^*(\mathbf{r}) d_{N\lambda}(\mathbf{r}')}{\sigma^2 + \Omega_{N\lambda}^2}, \quad (25)$$

where (24) uses the solutions of (18) to calculate the *bare* ($\lambda = 0$) response while (25) uses (19) to solve directly for the interacting response. The $\Re$ is unnecessary as the sum itself can be guaranteed real but may prove useful in some situations.

While $\hat{\chi}_\lambda$ defined by (25) has useful formal properties its direct evaluation may be numerically difficult and can be avoided. Unless otherwise noted we henceforth set $\mathbf{F}^{1\text{Int}} = \mathbf{0}$ and deal with internal interactions (when required) in a less direct, but precisely equivalent manner (as discussed tangentially in ref. 7 and in Appendix C of this manuscript).

IV. RELATIONSHIPS TO KS RESPONSE

In a Kohn-Sham system with orbitals $\psi_j(\mathbf{r}) \equiv \langle \mathbf{r} | j \rangle$, the exact bare response takes the form⁹

$$\chi_0^{\text{KS}}(\mathbf{r}, \mathbf{r}'; i\sigma) = -\Re \sum_{ai} \frac{\gamma_{ai}^*(\mathbf{r}) \gamma_{ai}(\mathbf{r}')}{\sigma^2 + \Omega_{ai}^2} \quad (26)$$

where γ_{ai} is a normalised transition density between unoccupied orbital $|a\rangle$ and occupied orbital $|i\rangle$ while Ω_{ai} is the transition frequency defined by

$$\gamma_{ai}(\mathbf{r}) = \sqrt{2\Omega_{ai}}\psi_a^*(\mathbf{r})\psi_i(\mathbf{r}), \quad (27a)$$

$$\Omega_{ai} = \epsilon_a - \epsilon_i > 0. \quad (27b)$$

As noted in Sec. I A, i is summed over occupied orbitals only and a over unoccupied orbitals only.

There is a transition current density associated with $|a\rangle$ and $|i\rangle$ which takes the form

$$\mathbf{j}_{ai}(\mathbf{r}) = \frac{1}{2i}[\psi_a(\mathbf{r})\nabla\psi_i^*(\mathbf{r}) - \psi_i^*(\mathbf{r})\nabla\psi_a(\mathbf{r})] \quad (28)$$

and where $i\nabla \cdot \mathbf{j}_{ai} = \Omega_{ai}\psi_i\psi_a^* = \sqrt{\Omega_{ai}/2}\gamma_{ai}$. Since $\{\mathbf{u}_N\}$ is complete (at least within a given finite basis) and orthonormal under (20) we can expand $\mathbf{j}_{ai}$ as

$$\mathbf{j}_{ai} = \sum_N K_{aiN} n^0 \mathbf{u}_N, \quad K_{aiN} = \int d\mathbf{r} \mathbf{u}_N^* \cdot \mathbf{j}_{ai}. \quad (29)$$

Taking the gradient of $\mathbf{j}_{ai}$ thus provides the following relationship between the KS transition densities γ_{ai} and the CM density modes d_N

$$\gamma_{ai} = \sum_N \frac{iK_{aiN}}{\sqrt{\Omega_{ai}/2}} d_N \quad (30)$$

so that any γ_{ai} can be expanded in $\{d_N\}$. Unfortunately the converse cannot be guaranteed except in the trivial one-electron case. In molecular systems we can make both d_N and ψ_j real so that

$$d_N = \sum_{ai} I_{Nai} \gamma_{ai} + \sum_{i < i'} I_{Ni'i} \gamma_{i'i} \quad (31)$$

where both i' and i are occupied while a is unoccupied.

Certain exact sum rules [equations (81)-(83) further discussed in appendix E of ref. 6] provide some further restrictions on the various coefficients. Since the f- and third-moment sum rules are satisfied by the CM it follows⁵ that

$$1 = \sum_{ai} h_{aiN}, \quad \Omega_N^2 = \sum_{ai} h_{aiN} \Omega_{ai}^2 \quad (32)$$

where Ω_{ai} are the Kohn-Sham transition frequencies if the CM equations involve the KS density n^0 and stress tensor $\mathbb{T}^0$. Here the *non-negative* weights h_{aiN} are defined via

$$h_{aiN} = \frac{2|K_{aiN}|^2}{\Omega_{ai}} \quad (33)$$

where K_{aiN} is defined in (29). More generally for $N \neq M$ these take the form

$$\sum_{ai} \frac{2K_{aiN}^* K_{aiM}}{\Omega_{ai}} = \delta_{NM}, \quad (34a)$$

$$\sum_{ai} \frac{2K_{aiN}^* K_{aiM}}{\Omega_{ai}} \Omega_{ai}^2 = \delta_{NM} \Omega_N^2 \quad (34b)$$

which come from inserting (30) into (26) and comparing the leading two powers of $1/\sigma^2$ with (24).

The second sum rule makes the relationship between Ω_N and the KS transition frequencies clear. We may also consider d_N to be an approximation to collections of the transition densities, with errors hopefully minimised by the sum rules and exact properties although no direct expansion exists. As discussed later these approximations become exact for one-electron (or $\uparrow\downarrow$) systems.

It is also worth noting that the lowest CM transition frequency Ω_1 can never be less than the transition frequency between the highest occupied- and lowest unoccupied- molecular orbital $\Omega_{LH} = \epsilon_L - \epsilon_H$. In the non-degenerate case the equality follows if and only if $h_{LH1} = 1$ with all other h_{ai1} zero. To prove the inequality we note that $\Omega_{ai} \geq \Omega_{LH}$ and thus

$$\Omega_1^2 \geq \sum_{ai} h_{ai1} \Omega_{LH}^2 \geq \Omega_{LH}^2. \quad (35)$$

If $h_{LH1} < 1$ then $h_{LH1} = 1 - \sum_{ai \neq LH} h_{ai1}$ and

$$\begin{aligned} \Omega_1^2 - \Omega_{LH}^2 &= (h_{LH1} - 1) \Omega_{LH}^2 + \sum_{ai \neq LH} h_{ai1} \Omega_{ai}^2 \\ &= \sum_{ai \neq LH} h_{ai1} (\Omega_{ai}^2 - \Omega_{LH}^2) > 0 \end{aligned}$$

since $h_{ai1} > 0$ and $\Omega_{ai}^2 - \Omega_{LH}^2 > 0$. Thus the equality only holds if $h_{ai1} = 1$.

V. CORRELATION ENERGIES MADE SIMPLE

Once we have obtained d_N and Ω_N , the CM approximation to the correlation energy can be calculated.⁷ We define the Coulomb projection matrix $\mathbb{W}$ with elements

$$W_{NM} = \int d\mathbf{r} \mathbf{u}_M^*(\mathbf{r}) \cdot [\hat{\mathbf{Q}} \mathbf{u}_N](\mathbf{r}) \quad (36)$$

$$= \int d\mathbf{r} d\mathbf{r}' \frac{d_N(\mathbf{r}) d_M^*(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} \quad (37)$$

and $\mathbb{L}$ with elements $L_{MN} = \delta_{MN}\Omega_N^2$. Through the working in Appendix C we can show that the correlation energy is

$$E_c = \int_0^1 d\lambda \int_0^\infty \frac{d\sigma}{\pi} \times \frac{1}{2} \text{Tr} \left[\frac{\mathbb{W}}{\sigma^2 + \mathbb{L} + \lambda \mathbb{W}} - \frac{\mathbb{W}}{\sigma^2 + \mathbb{L}} \right] \quad (38)$$

or we can use the Furche-like²⁵ integrated form

$$E_c = \frac{1}{2} \sum_N \left[\bar{\Omega}_N - \Omega_N \left(1 + \frac{W_{NN}}{2\Omega_N^2} \right) \right] \quad (39)$$

where $\bar{\Omega}_N^2$ are the eigenvalues of $\hat{\mathbf{R}} + \hat{\mathbf{Q}}$ or $\mathbb{L} + \mathbb{W}$. As discussed in Appendix C the two diagonalisations are formally equivalent but experience in similar techniques suggests that working in the transition densities of the bare response will allow for better convergence. In practise diagonalising $\mathbb{L} + \mathbb{W}$ is expected to be faster and numerically more reliable and robust.

Using the eigenvalues of $\mathbb{L} + \mathbb{W}$ has a further advantage: we can use a perturbative solution to find the eigenvalues of $\mathbb{L} + \mathbb{W}$ if $\Omega_N^2 \gg \bar{W}_N$ where $\bar{W}_N = \sum_M |W_{NM}|$ (see Appendix C for details). We define an N^* such that $\Omega_N^2 \geq K \bar{W}_N \forall N > N^*$ where K is sufficiently large. We then solve the reduced $N^* \times N^*$ eigen-equation $\mathbb{L}^* + \mathbb{W}^*$ to obtain $\bar{\Omega}_{N=1\dots N^*}^*$ and calculate

$$E_c \approx \frac{1}{2} \sum_{N=1}^{N^*} [\bar{\Omega}_N^* - \Omega_N (1 + \beta_N)] - \sum_{N > N^*} \frac{\Omega_N \beta_N^2}{4} \quad (40)$$

where $\beta_N = \frac{W_{NN}}{2\Omega_N^2}$.

VI. ORTHONORMAL DISPLACEMENT

If we consider the orthonormality condition (20) of the CM eigenmodes in a bound system we can see that $\mathbf{u}_N$ may be permitted to grow as $|\mathbf{r}| \rightarrow \infty$ provided $\sqrt{n^0} \mathbf{u}_N$ decreases. In atomic and molecular systems all valid solutions will, in fact, grow exponentially due to the asymptotic form of the orbitals. This suggests that an orthonormal fluid displacement $\boldsymbol{\xi} = \sqrt{n^0} \mathbf{u}$ will be a more natural quantity to use in these systems as it is guaranteed to decrease. Here the orthonormal modes are

$$\boldsymbol{\xi}_N = \sqrt{n^0} \mathbf{u}_N \quad (41)$$

where $\int d\mathbf{r} \boldsymbol{\xi}_N \cdot \boldsymbol{\xi}_M = \delta_{NM}$

While $\mathbf{u} = \sum_N c_N \mathbf{u}_N$ has a well-defined physical meaning (the fluid displacement), $\boldsymbol{\xi} = \sum c_N \boldsymbol{\xi}_N$ is somewhat harder to interpret. However some insight can be gained if we define the groundstate quasi-orbital $\Psi^0 = \sqrt{n^0}$ and its first order change Ψ^1 . Since $n^1 = 2\Psi^0\Psi^1 = -\nabla \cdot \Psi^0 \boldsymbol{\xi}$ it is clear that

$$2\Psi^1 = - \left(\nabla + \frac{[\nabla \Psi^0]}{\Psi^0} \right) \cdot \boldsymbol{\xi} \quad (42)$$

and thus $\boldsymbol{\xi}$ is related to the perturbation of the quasi-orbital. It is also related, via (6), to the groundstate properties n^1 and $\mathbf{j}$ through

$$\mathbf{j} = \partial_t \Psi^0 \boldsymbol{\xi} \quad n^1 = -\nabla \Psi^0 \boldsymbol{\xi}. \quad (43)$$

We must calculate $\boldsymbol{\xi}_N$ through the CM equations (18)

$$\Omega_N^2 \boldsymbol{\xi}_N = \hat{R}^{(\epsilon)} \boldsymbol{\xi}_N, \quad (44)$$

$$\hat{R}^{(\epsilon)} = \frac{1}{\Psi^0} \hat{R} \frac{1}{\Psi^0} \quad (45)$$

where $\hat{R}$ is defined in equation (11). The change in density of a given mode becomes

$$d_N(\mathbf{r}) = -\Psi^0 (\nabla + \boldsymbol{\eta}) \cdot \boldsymbol{\xi}_N \quad (46)$$

where

$$\boldsymbol{\eta} = \frac{[\nabla n^0]}{2n^0} = \frac{[\nabla \Psi^0]}{\Psi^0} = \nabla \log \Psi^0 = \frac{1}{2} \nabla \log n^0 \quad (47)$$

is the logarithmic gradient of Ψ^0 . Equation 46 can be used in (37) to calculate the matrix elements of $\mathbb{W}$ for use in correlation energy calculations

Inserting (11) into (45) gives

$$\begin{aligned} \hat{R}_{\mu\nu}^{(\epsilon)} = & -\frac{1}{\Psi^0} (\partial_\alpha T_{\mu\nu}^0 \partial_\alpha + \partial_\alpha T_{\alpha\nu}^0 \partial_\mu + \partial_\nu T_{\alpha\mu}^0 \partial_\alpha) \frac{1}{\Psi^0} \\ & + V_{,\mu\nu}^{\text{KS}} + \frac{1}{4} \frac{1}{\Psi^0} \partial_\nu \partial_\alpha n \partial_\alpha \partial_\mu \frac{1}{\Psi^0}. \end{aligned} \quad (48)$$

Using the derivative operator identity

$$\frac{1}{\Psi^0} \partial_\alpha - \partial_\alpha \frac{1}{\Psi^0} = \frac{\eta_\alpha}{\Psi^0}. \quad (49)$$

and defining $t_{\mu\nu} = T_{\mu\nu}^0/n^0$, allows us to convert (48) into the following succinct reformulation

$$\begin{aligned}\hat{R}^{(\boldsymbol{\xi})} = & V_{,\mu\nu}^{\text{KS}} + \frac{1}{4}(\partial_\nu + \eta_\nu)(\nabla^2 - S)(\partial_\mu - \eta_\mu) \\ & - (\partial_\alpha + \eta_\alpha)t_{\mu\nu}(\partial_\alpha - \eta_\alpha) \\ & - (\partial_\alpha + \eta_\alpha)t_{\alpha\nu}(\partial_\mu - \eta_\mu) \\ & - (\partial_\nu + \eta_\nu)t_{\alpha\mu}(\partial_\alpha - \eta_\alpha)\end{aligned}\quad (50)$$

which we can use in (44). Here

$$\begin{aligned}S = & \nabla^2 - (\nabla + \boldsymbol{\eta}) \cdot (\nabla - \boldsymbol{\eta}) \\ = & [\partial_\alpha \eta_\alpha] + \eta_\alpha \eta_\alpha = ([\frac{1}{2}\nabla^2 n^0]/n^0 - \eta_\alpha \eta_\alpha).\end{aligned}\quad (51)$$

All functions appearing in these expressions depend on groundstate orbital wavefunctions and their derivatives only. Expanding in terms of occupied orbitals they are

$$t_{\mu\nu} = \frac{\Re \frac{1}{2} \sum_i ([\partial_\mu \psi_i^*][\partial_\nu \psi_i] - \psi_i^* [\partial_\mu \partial_\nu \psi_i])}{\sum_i \psi_i^* \psi_i}, \quad (52)$$

$$\eta_\alpha = \frac{\Re \sum_i \psi_i^* [\partial_\alpha \psi_i]}{\sum_i \psi_i^* \psi_i}, \quad (53)$$

$$S = \frac{\Re \sum_i ([\partial_\alpha \psi_i^*][\partial_\alpha \psi_i] + \psi_i^* [\nabla^2 \psi_i])}{\sum_i \psi_i^* \psi_i} - \eta_\alpha \eta_\alpha. \quad (54)$$

Here the force balance equation $V_{,\mu}^{\text{KS}} = -(\partial_\alpha + 2\eta_\alpha)t_{\alpha\mu}$ replaces the usual $n^0 V_{,\mu}^{\text{KS}} = -\partial_\alpha T_{\alpha\mu}^0$.

In a one-electron system with occupied orbital ψ and energy ϵ_0 these reduce to

$$\eta_\alpha = \frac{\psi_{,\alpha}}{\psi}, \quad t_{\mu\nu} = \frac{1}{2}(\eta_\mu \eta_\nu - \frac{\psi_{,\mu\nu}}{\psi}), \quad (55a)$$

$$S = 2(V^{\text{KS}} - \epsilon_0). \quad (55b)$$

It is worth noting that, for molecular systems with Coulomb-like nuclear potentials the outermost tail is dominated by one-electron-like behaviour and:

1. The denominators of $t_{\mu\nu}$, η_α and S are densities and thus everywhere positive,
2. $\lim_{r \rightarrow \infty} t_{\mu\nu}(\mathbf{r}) \sim 0$,
3. $\lim_{r \rightarrow \infty} |\boldsymbol{\eta}(\mathbf{r})| \sim |\sqrt{-2\epsilon_H}|$ where ϵ_H is the KS eigenvalue of the highest occupied orbital,

$$4. \lim_{r \rightarrow \infty} S(\mathbf{r}) \sim 2(V^{\text{KS}}(\mathbf{r}) - \epsilon_H).$$

where $\mathbf{r}$ is the displacement from the center of the highest occupied orbital and we ignore leading terms that decay exponentially. In large molecules these expressions may also hold true closer to nuclei A with ϵ_H replaced by the local ϵ_H^A .

VII. ONE-ELECTRON SYSTEMS

Following equations (41) and (45-46) of ref. 4, it is possible to show that the non-linearised CM formalism is equivalent to Madelung hydrodynamics¹ in a Lagrangian frame for one electron systems. If all external fields involve gradients of scalar potentials only, this is directly equivalent²⁶ to finding solutions of the one-electron Schrödinger equation (SE). Here we show this equivalence directly from the SE in the linear response limit required by the density response $\hat{\chi}_0$.

We proceed to prove this as follows: i) we derive the relationship between the perturbed one-electron wavefunction and its “orthonormal displacement vector” $\boldsymbol{\xi}^{1e^-}$ to show that the latter is entirely determined by the former (and vice versa up to a trivial phase via the Runge-Gross theorem⁹); ii) we show that finding a free-standing solution of the linear-perturbed SE for a one-electron system is equivalent to solving an equation of form $\omega^2 \boldsymbol{\xi}^{1e^-} = \hat{\mathbf{R}}^{1e^-} \boldsymbol{\xi}^{1e^-}$; and iii) we show that $\hat{\mathbf{R}}^{1e^-} = \hat{\mathbf{R}}$ as defined in (11).

In a one-electron system we can set $V = V^{\text{KS}} - \epsilon_0$ to eliminate the KS energy of the single occupied orbital. Thus the groundstate Hamiltonian takes the form

$$\left(-\frac{1}{2}\nabla^2 + V\right)\psi \equiv \hat{h}\psi = 0 \quad (56)$$

where ψ is the only occupied electron wavefunction which we make real. If we apply a small, time-dependent external potential $V^1(t)$ then we can find a new solution $\psi'(t)$ through

$$[\hat{h} + V^1(t)]\psi'(t) = -i\partial_t\psi'(t). \quad (57)$$

which will be perturbed only slightly from the groundstate solution. We can write the perturbed wavefunction via a change to its magnitude and a rotation of its phase such that

$$\psi'(t) = [\psi + \psi^1(t)]e^{i\phi^1(t)} \quad (58)$$

or, truncating to first order, $\psi'(t) \approx \psi + \psi^1(t) + i\psi\phi^1(t)$. Thus it is sufficient to calculate ϕ^1 and ψ^1 to fully determine the perturbed solution.

Inserting (58) into (57) and matching real and imaginary components (to linear order) gives

$$\psi\partial_t\phi^1(t) = \psi V^1(t) + \hat{h}\psi^1(t), \quad (59)$$

$$\partial_t\psi^1(t) = (\nabla\psi) \cdot [\nabla\phi^1(t)] + \frac{1}{2}\psi\nabla^2\phi^1(t). \quad (60)$$

where the second expression relates $\partial_t\psi^1$ directly to $\nabla\phi^1$. If we then assume a time-periodic external potential $V^1(t) = V^1 e^{i\omega t}$ it follows that $\psi^1(t) = \psi^1 e^{i\omega t}$ and $\phi^1(t) = \phi^1 e^{i\omega t}$ and thus $\partial_t \equiv i\omega$. We can then use (60) to eliminate ψ^1 and derive the equations governing density perturbations in terms of ϕ^1 only. Here

$$-\omega^2\phi^1 = \frac{1}{2}\psi^{-1}\hat{h}(\psi_{,\alpha} + \partial_\alpha\psi)\partial_\alpha\phi^1 + i\omega V^1(t), \quad (61)$$

$$n^1 = 2\psi\psi^1 = -\frac{\psi}{-i\omega}(\psi_{,\alpha} + \partial_\alpha\psi)\partial_\alpha\phi^1 \quad (62)$$

where n^1 is the linear change in density and we have used the derivative relationship $\partial_\alpha\psi - \psi\partial_\alpha \equiv \psi_{,\alpha}$. By the Runge-Gross theorem⁹ $V' = V + V^1(t)$ is a functional of the density $n' = n^0 + n^1(t)$ only in the linear response regime, and from (62) it is clear that ψ and $\nabla\phi^1$ are sufficient to determine electronic properties.

The Schrödinger current density $\mathbf{j}$ of the perturbed, one-electron system is calculated through

$$\mathbf{j}^{1e^-} = \frac{1}{2i}[\psi'\nabla\psi'^* - \psi'^*\nabla\psi'] \approx -\psi^2\nabla\phi^1 \quad (63)$$

where we use the first-order perturbation expression $\psi' \approx \psi + \psi^1 + i\psi\phi^1$ to derive the second identity. In general²⁴ the displacement $\mathbf{u}$ is related to the current via (6) and thus $i\omega\mathbf{u}^{1e^-} = -\nabla\phi^1$ for the one-electron system.

It then follows trivially that the normalised displacement $\boldsymbol{\xi}^{1e^-} = \sqrt{n^0}\mathbf{u}^{1e^-} \equiv \psi\mathbf{u}^{1e^-}$ is related to ϕ^1 via

$$\boldsymbol{\xi}^{1e^-} \equiv \frac{\psi}{-i\omega}\nabla\phi^1 \quad (64)$$

and, from the Runge-Gross theorem,⁹ that ϕ^1 can be obtained from $\boldsymbol{\xi}^{1e^-}$. Using (62) the density perturbation takes the expected form (43) $n^1 = -\nabla \cdot \psi\boldsymbol{\xi}^{1e^-}$. This completes the first stage of the proof.

By taking the gradient of (61) and using $\nabla\phi^1 = (-i\omega/\psi)\xi^{1e^-}$ [from (64)] we can explicitly solve the free-standing ($V^1 = 0$) equation for ξ^{1e^-} via

$$-\omega^2 \frac{\xi_\mu^{1e^-}}{\psi} = \frac{1}{2} \partial_\mu \psi^{-1} \hat{h}(\psi_{,\nu} + \partial_\nu \psi) \frac{\xi_\nu^{1e^-}}{\psi} \quad (65)$$

$$\omega^2 \xi_\mu^{1e^-} = \frac{1}{4} (\partial_\mu - \eta_\mu) (\nabla^2 - S) (\partial_\nu + \eta_\nu) \xi_\nu^{1e^-} \quad (66)$$

where we used the one-electron specific relationships (55) $\eta_\alpha = \psi_{,\alpha}/\psi$ and $2\hat{h} = S - \nabla^2$ for the second expression. Thus we can define a linear operator

$$\hat{R}_{\mu\nu}^{1e^-} = \frac{1}{4} (\partial_\mu - \eta_\mu) (\nabla^2 - S) (\partial_\nu + \eta_\nu) \quad (67)$$

such that $\omega^2 \xi_\mu^{1e^-} = \hat{R}_{\mu\nu}^{1e^-} \xi_\nu^{1e^-}$ which is of the same form as (8), as desired for the second stage of the proof.

Finally it remains to be shown that $\hat{R} = \hat{R}^{1e^-}$. Following the working in Appendix D we show that

$$\begin{aligned} \hat{R}_{\mu\nu}^{1e^-} &= V_{,\mu\nu} + \frac{1}{4} (\partial_\nu + \eta_\nu) (\nabla^2 - S) (\partial_\mu - \eta_\mu) \\ &\quad - (\partial_\alpha + \eta_\alpha) t_{\mu\nu} (\partial_\alpha - \eta_\alpha) \\ &\quad - (\partial_\alpha + \eta_\alpha) t_{\alpha\nu} (\partial_\mu - \eta_\mu) \\ &\quad - (\partial_\nu + \eta_\nu) t_{\alpha\mu} (\partial_\alpha - \eta_\alpha) \\ &\equiv \hat{R}_{\mu\nu}^{(\xi)} \end{aligned} \quad (68)$$

and the proof is complete.

Thus we have shown that i) ξ^{1e^-} is bijectively (up to a phase) related to the first-order solution of the perturbed SE; ii) It obeys $\omega^2 \xi^{1e^-} = \hat{R}^{1e^-} \xi^{1e^-}$ for free-standing modes and iii) $\hat{R}^{1e^-} = \hat{R}$. Thus the governing equation is identical in both cases and it follows that $\xi^{1e^-} \equiv \xi$. This confirms that the solutions of the CM equations are directly equivalent to the solutions of the perturbed Schrödinger equation in one-electron systems, and that the CM is thus exact.

A. Transition modes of one-electron systems

In a one-electron system, the density response $\hat{\chi}_0$ calculated via orbital transitions (26) must be the same as that calculated via the CM transitions (24). Thus

$$\chi_0 = \sum_a \frac{\gamma_{a0}(\mathbf{r})\gamma_{a0}(\mathbf{r}')}{\Omega_{a0}^2 + \sigma^2} = \sum_N \frac{d_N(\mathbf{r})d_N(\mathbf{r}')}{\Omega_N^2 + \sigma^2}. \quad (69)$$

where a is summed over the unoccupied orbitals, while $\Omega_{a0} = \epsilon_a - \epsilon_0$ and $\gamma_{a0} = \pm\sqrt{2\Omega_{a0}}\psi\psi_a$. Since the equality must be true for all σ it follows that the individual numerators and denominators of the sums must be paired and we can choose an N_a such that $\Omega_{N_a} = \Omega_{a0}$ and $d_{N_a}(\mathbf{r}) = \gamma_{a0}(\mathbf{r})$.

Equating (46) and (27) gives

$$d_{N_a} = -\psi(\nabla + \boldsymbol{\eta}) \cdot \boldsymbol{\xi}_{N_a} = \sqrt{2\Omega_{a0}}\psi\psi_a. \quad (70)$$

This has a (non-unique) solution

$$\boldsymbol{\xi}_{N_a} = \frac{1}{\sqrt{2\Omega_{a0}}}(\nabla - \boldsymbol{\eta})\psi_a \quad (71)$$

which must therefore be a valid solution of the CM equations. The constant pre-factor comes from $-(\nabla + \boldsymbol{\eta}) \cdot (\nabla - \boldsymbol{\eta}) \equiv 2(\hat{h} - \epsilon_0)$ in a one-electron system.

Tao *et al*⁵ tested the one-electron exactness on general s -transitions in the Hydrogen atom. We illustrate (70) and (71) on the $1s$ to $2p$ transition. Here the $1s$ orbital is occupied with orbital $\psi_{1s} = e^{-r}/\sqrt{\pi}$ and energy $\epsilon_0 = \epsilon_{1s} = -\frac{1}{2}$ while the $2p_z$ orbital has $\psi_{2p_z} = ze^{-r/2}/\sqrt{32\pi}$ and $\epsilon_{2p_z} = -\frac{1}{8}$. The density is thus $n^0 = e^{-2r}/\pi$ and $\boldsymbol{\eta} = -\hat{\mathbf{r}}$.

Using $\nabla ze^{-r/2} = (\hat{\mathbf{z}} - \frac{1}{2}z\hat{\mathbf{r}})e^{-r/2}$ we can test (70) and (71). With $\Omega_{2p_z-1s} = \frac{3}{8}$ we find

$$\boldsymbol{\xi}_{2p_z-1s} = \frac{1}{\sqrt{96\pi}}e^{-r/2}(z\hat{\mathbf{r}} + 2\hat{\mathbf{z}}), \quad (72)$$

$$\begin{aligned} d_{2p_z-1s} &= -\frac{\psi_{1s}}{\sqrt{96\pi}}(\nabla - \hat{\mathbf{r}}) \cdot (z\hat{\mathbf{r}} + 2\hat{\mathbf{z}})e^{-r/2} \\ &= \sqrt{\frac{3}{4}}\psi_{1s}\psi_{2p_z}. \end{aligned} \quad (73)$$

in agreement with (70) and where $3/4 = 2\Omega_{1s-2p_z}$ as expected. Note that the usual displacement vector $\mathbf{u}_{2p_z-1s} = \frac{e^{r/2}}{\sqrt{96}}(z\hat{\mathbf{r}} + 2\hat{\mathbf{z}})$ grows exponentially with r .

VIII. SCALAR APPROXIMATION TO CM

In many-electron systems, equation (71) is no longer guaranteed to be true. However, as we can write any well-behaved vector function as a gradient plus a curl we may set

$$\boldsymbol{\xi}_N(\mathbf{r}) = \Psi^0 \{ \nabla s_N(\mathbf{r}) - \nabla \times \mathbf{v}_N(\mathbf{r}) \} \quad (74)$$

where $s_N(\mathbf{r})$ and $\mathbf{v}_N(\mathbf{r})$ are arbitrary scalar/vector functions with appropriate asymptotes (we choose our gauge to ensure $\nabla \cdot \mathbf{v}_N = 0$). The form of (71) suggests that we might approximate our eigen-solutions $\boldsymbol{\xi}_N$ by setting

$$\tilde{\boldsymbol{\xi}}_N(\mathbf{r}) \approx \{ \nabla - \boldsymbol{\eta}(\mathbf{r}) \} \phi_N(\mathbf{r}) = \Psi^0(\mathbf{r}) \nabla \frac{\phi_N(\mathbf{r})}{\Psi^0(\mathbf{r})} \quad (75)$$

which is equivalent to making the approximation $s_N \approx \phi_N/\Psi^0$ and $\mathbf{v}_N \approx \mathbf{0}$. This approximation will be exact for one-electron systems and one-dimensional (1D) systems²⁷ but is not true in general.

The regular CM eigen-equation (44) is equivalent to finding stationary solutions $\delta\mathcal{E}/\delta\boldsymbol{\xi} = \mathbf{0}$ of $\mathcal{E}[\boldsymbol{\xi}] = \frac{1}{2} \int d\mathbf{r} \boldsymbol{\xi}^* \cdot \hat{\mathbf{R}} \cdot \boldsymbol{\xi}$ for vectors $\boldsymbol{\xi}$ satisfying $\int d\mathbf{r} \boldsymbol{\xi}^* \cdot \boldsymbol{\xi} = 1$. We called these solutions $\boldsymbol{\xi}_N$ and they can be found through $\Omega_N^2 \boldsymbol{\xi}_N = \hat{\mathbf{R}} \boldsymbol{\xi}_N$ where $\hat{\mathbf{R}}$ is defined in (11). Under the scalar approximation we restrict our solutions to vectors $\tilde{\boldsymbol{\xi}}$ expressible as $(\nabla - \boldsymbol{\eta})\phi$ which form a connected subspace of all possible $\boldsymbol{\xi}$. We thus look for solutions $\tilde{\boldsymbol{\xi}}_N(\phi)$ of $\mathcal{E}[\boldsymbol{\xi}]$ which are stationary under variation of ϕ ie. $\delta\mathcal{E}[\boldsymbol{\xi}(\phi)]/\delta\phi = 0$ subject to $\int d\mathbf{r} \tilde{\boldsymbol{\xi}}(\phi) \cdot \tilde{\boldsymbol{\xi}}(\phi) = 1$.

The restricted solutions can be found directly by setting $\tilde{\mathcal{E}}[\phi] = -\frac{1}{2} \int d\mathbf{r} \phi^* (\nabla + \boldsymbol{\eta}) \cdot \hat{\mathbf{R}} \cdot (\nabla - \boldsymbol{\eta})\phi$ and setting the constraint to $-\int d\mathbf{r} \phi^* (\nabla + \boldsymbol{\eta}) \cdot (\nabla - \boldsymbol{\eta})\phi = 1$. The general Hermitian eigen-equation for ϕ_N thus becomes

$$\tilde{\Omega}_N^2 \hat{N}^\phi \phi_N = \hat{R}^\phi \phi_N \quad (76)$$

with orthogonal solutions normalised under $\int d\mathbf{r} \phi_N^* \hat{N}^\phi \phi_M = \delta_{NM}$. Here

$$\hat{N}^\phi = -(\nabla + \boldsymbol{\eta}) \cdot (\nabla - \boldsymbol{\eta}) = S - \nabla^2 \quad (77)$$

$$\hat{R}^\phi = -(\partial_\mu + \eta_\mu) \hat{R}_{\mu\nu} (\partial_\nu - \eta_\nu). \quad (78)$$

It is obvious that $\tilde{\Omega}_1^2 \geq \Omega_1^2$ since the subspace minimum of $\mathcal{E}$ must be equal to or higher than its true minimum.

Such an approximation loses some accuracy and some nice properties of the true CM but reduces the problem from a tensor to a scalar. Its exactness in a variety of systems including

the one-electron case in any number of dimensions suggests that it might be appropriate for vdW calculations in molecular systems. Here the vdW physics are often dominated by the asymptotic regions which show one-electron-like behaviour (see Section VIII C for further details).

A. Scalar approximation in KS orbitals

We can expand $\phi_N = \sum_j p_{Nj} \psi_j$ in the KS orbitals (or any other complete and orthonormal basis set) so that, from (75),

$$\tilde{\boldsymbol{\xi}}_N(\mathbf{r}) = \sum_j p_{Nj} \{\nabla - \boldsymbol{\eta}(\mathbf{r})\} \psi_j(\mathbf{r}) \quad (79)$$

where ψ_j are KS orbitals (occupied or otherwise). In reality we must truncate to the lowest N_{Bas} orbitals. The transition density modes are thus (remembering that $\sqrt{n^0} = \Psi^0$)

$$\tilde{d}_N(\mathbf{r}) = -\Psi^0 \sum_j p_{Nj} (\nabla + \boldsymbol{\eta}) \cdot (\nabla - \boldsymbol{\eta}) \psi_j \quad (80a)$$

$$= \Psi^0 \sum_j p_{Nj} (S - \nabla^2) \psi_j \quad (80b)$$

$$= \Psi^0 \sum_j p_{Nj} [2(\epsilon_k - V^{\text{KS}}) + S] \psi_j. \quad (80c)$$

where the first two properties are true for any basis set but the third property is only true for the KS orbitals.

Projection into the KS orbitals allows some insight into the physical meaning of this approximation to be obtained by considering the quasi-orbital Ψ^0 . Here we can set $\tilde{\boldsymbol{\xi}}_N = (2i/\Psi^0) \sum_j p_{Nj} \mathbf{j}_j$ where the quasi-transition current

$$\mathbf{j}_j = \frac{1}{2i} [\Psi^0 (\nabla - \boldsymbol{\eta}) \psi_j] = \frac{1}{2i} [\Psi^0 \nabla \psi_j - \psi_j \nabla \Psi^0] \quad (81)$$

has a similar form to a transition current $\mathbf{j}_{ai} = \frac{1}{2i} (\psi_i \nabla \psi_a - \psi_a \nabla \psi_i)$ with the occupied orbital replaced by the quasi-orbital of the *total* density.

If we pre-multiply (44) by $\tilde{\boldsymbol{\xi}}_M$ and integrate we find $\tilde{\Omega}_N^2 \int d\mathbf{r} \tilde{\boldsymbol{\xi}}_M \cdot \tilde{\boldsymbol{\xi}}_N = \int d\mathbf{r} \tilde{\boldsymbol{\xi}}_M \hat{R} \tilde{\boldsymbol{\xi}}_N$ subject to $\int d\mathbf{r} \tilde{\boldsymbol{\xi}}_M \cdot \tilde{\boldsymbol{\xi}}_N = \delta_{MN}$. Using (79) the two integrals become

$$\int d\mathbf{r} \tilde{\boldsymbol{\xi}}_M^* \cdot \tilde{\boldsymbol{\xi}}_N = \sum_{jk} p_{Mj}^* p_{Nk} N_{jk}, \quad (82)$$

$$\int d\mathbf{r} \tilde{\boldsymbol{\xi}}_M^* \hat{R} \tilde{\boldsymbol{\xi}}_N = \sum_{jk} p_{Mj}^* p_{Nk} R_{jk}. \quad (83)$$

where

$$N_{jk} = \int d\mathbf{r} [(\partial_\alpha - \eta_\alpha)\psi_j^*][(\partial_\alpha - \eta_\alpha)\psi_k], \quad (84)$$

$$R_{jk} = \int d\mathbf{r} [(\partial_\mu - \eta_\mu)\psi_j^*]\hat{R}_{\mu\nu}[(\partial_\nu - \eta_\nu)\psi_k]. \quad (85)$$

Taking derivatives with respect to p_{Mj} , the eigen-equation (44) thus becomes

$$\tilde{\Omega}_N^2 N_{jk} p_{Nk} = R_{jk} p_{Nk} \quad (86)$$

subject to the orthogonality condition $\sum_{jk} N_{jk} p_{Mj}^* p_{Nk} = \delta_{NM}$. Following the details of Appendix E we find

$$\begin{aligned} R_{jk} = & \int d\mathbf{r} [\hat{D}_\mu \psi_j^*] V_{,\mu\nu} [\hat{D}_\nu \psi_k] \\ & + \frac{1}{4} \int d\mathbf{r} [\hat{D}_\alpha \hat{D}_\mu \hat{D}_\nu \psi_j^*] [\hat{D}_\alpha \hat{D}_\mu \hat{D}_\nu \psi_k] \\ & + 3 \int d\mathbf{r} [\hat{D}_\alpha \hat{D}_\mu \psi_j^*] t_{\mu\nu} [\hat{D}_\alpha \hat{D}_\nu \psi_k] \end{aligned} \quad (87)$$

where $\hat{D}_\alpha \equiv \partial_\alpha - \eta_\alpha$.

B. Matrix form of the scalar approximation

Since all terms involve repeated use of the operator $(\partial_\alpha - \eta_\alpha)$ we may simplify things somewhat by adopting a matrix notation to represent this operator. For a finite basis set of size N_{Bas} we define the following $N_{\text{Bas}} \times N_{\text{Bas}}$ matrices: $\mathbb{D}_\alpha$, $\mathbb{T}_{\mu\nu}$ and $\mathbb{V}_{\mu\nu}$ with elements

$$D_{\alpha jk} = \int d\mathbf{r} \psi_j^* (\partial_\alpha - \eta_\alpha) \psi_k, \quad (88)$$

$$T_{\mu\nu jk} = \int d\mathbf{r} \psi_j^* t_{\mu\nu} \psi_k, \quad (89)$$

$$V_{\mu\nu jk} = \int d\mathbf{r} \psi_j^* V_{,\mu\nu}^{\text{KS}} \psi_k. \quad (90)$$

We can also integrate by parts to obtain $V_{\mu\nu jk} = \int d\mathbf{r} V^{\text{KS}} [\partial_\mu \partial_\nu \psi_j^* \psi_k]$. Noting that the orbitals ψ_j form an orthonormal set we see that $(\partial_\alpha - \eta_\alpha)\psi_k = \sum D_{\alpha jk} \psi_j$ with similar relationships for the others.

This allows us to write the matrix eigen-equation

$$\tilde{\Omega}_N^2 \mathbb{N} \mathbb{P}_N = \mathbb{R} \mathbb{P}_N \quad (91)$$

where $\mathbb{P}_N$ is an $N_{\text{Bas}} \times 1$ matrix with elements p_{Nj} . Here

$$\mathbb{N} = \mathbb{D}_\alpha^\dagger \mathbb{D}_\alpha, \quad (92)$$

$$\mathbb{R} = \mathbb{D}_\mu^\dagger \mathbb{V}_{\mu\nu} \mathbb{D}_\nu + \mathbb{D}_\mu^\dagger \mathbb{D}_\alpha^\dagger \left(3\mathbb{T}_{\mu\nu} + \frac{1}{4}\mathbb{D}_\nu^\dagger \mathbb{D}_\mu \right) \mathbb{D}_\alpha \mathbb{D}_\nu. \quad (93)$$

Orthogonality is given by $\mathbb{P}_N^\dagger \mathbb{N} \mathbb{P}_M = \delta_{NM}$. We note that all tensor components are bundled into $\mathbb{N}$ and $\mathbb{R}$ via summation over α , μ and ν . These equations are only true in the strict limit of an infinite number of orbitals but will converge to the true approximate solution in typical systems.

C. One-electron like asymptotic behaviour

In the asymptotic $|\mathbf{r}| \rightarrow \infty$ region of a localised system, the highest occupied molecular orbital (HOMO) ψ_H with occupation number f_H dominates [at least in the non-degenerate case or for degenerate system with spherical symmetry where we set $f_H = (2l_H + 1)$ and $\psi_H(\mathbf{r}) = R_{n_H}(r)$, the radial component of $\psi_{n_H l_H m}(\mathbf{r}) = R_{n_H}(r) Y_{l_H m}(\Omega_r)$]. Thus, in the limit $\mathbf{r} \rightarrow \infty$ we can set:

$$n^0 \approx f_H |\psi_H|^2, \quad \boldsymbol{\eta} \approx \frac{[\nabla \psi_H]}{\psi_H}, \quad (94)$$

$$t_{\mu\nu} \approx -\frac{1}{2} \partial_\mu \eta_\nu, \quad S \approx 2(V^{\text{KS}} - \epsilon_H). \quad (95)$$

All equations reduce to the one-electron form in these limits, with the single occupied orbital given by the HOMO.

We can check how close a region of a many-electron system is to a one-electron-like system by considering the difference of S from its asymptotic form. Here we set

$$\tilde{S}(\mathbf{r}) = S(\mathbf{r}) - 2(V^{\text{KS}}(\mathbf{r}) - \epsilon_H) \quad (96a)$$

$$= C^0(\mathbf{r}) + 2(\epsilon_H - \tilde{\epsilon}(\mathbf{r})) \quad (96b)$$

where

$$C^0(\mathbf{r}) = \sum_i \frac{f_i |\nabla \psi_i(\mathbf{r})|^2}{n^0(\mathbf{r})} - |\boldsymbol{\eta}(\mathbf{r})|^2 \quad (97)$$

$$\tilde{\epsilon}(\mathbf{r}) = \sum_i \frac{f_i |\psi_i(\mathbf{r})|^2}{n^0(\mathbf{r})} \epsilon_i \quad (98)$$

and where C^0 is related to the Fermi-hole curvature²⁸ of the KS system. In the limit $|\mathbf{r}| \rightarrow \infty$ it is clear that $C^0(\mathbf{r}) = 0$ and $\tilde{\epsilon}(\mathbf{r}) = \epsilon_H$ (true everywhere for one-electron systems).

This outer region is crucial for van der Waals physics in many systems. Typically only transitions within a small frequency range, especially that to the lowest unoccupied molecular orbital (LUMO), will dominate in this region. In such transitions $\int d\mathbf{r} \psi_k^* \tilde{S} \psi_k$ will be small. The asymptotic ‘exactness’ in this limit suggest strongly that the scalar approximation to CM is appropriate for calculating vdW forces.

Even a well chosen but simple approximation, using a limited number of unoccupied orbitals and quasi-orbitals, may provide quite accurate estimations of the vdW physics. The ability of the CM (and the scalar approximation) to make collective modes for the transition physics may aid in convergence compared to full dRPA calculations.

D. Transitions and energy calculations in the scalar approximation

Once we have obtained solutions of (91) we can use p_{Nj} to evaluate d_N through (80). Using $\tilde{\Omega}_N$ also we can then evaluate $\hat{\chi}_0$ and properties which depend on it such as the correlation energy.

We can also use the asymptotic form of $S(\mathbf{r})$ to further investigate d_N and through it correlations. Combining (80) and (96) lets us write

$$\begin{aligned} d_N &= \Psi^0 \sum_j p_{Nj} \left[2(V^{\text{KS}} - \epsilon_H) - \nabla^2 + \tilde{S} \right] \psi_j \\ &= \Psi^0 \sum_j p_{Nj} \left(2\Omega_{jH} + \tilde{S} \right) \psi_j \end{aligned} \quad (99)$$

where the $\Omega_{jH} = \epsilon_j - \epsilon_H$ comes from applying the Schrödinger Hamiltonian $-\frac{1}{2}\nabla^2 + V^{\text{KS}}$ to ψ_j . Here $\tilde{S}$ covers the deviation of many-electron systems from their one-electron equivalents.

Let us expand (99) as follows:

$$d_N = \Psi^0 \sum_j d_{Nj} \psi_j. \quad (100)$$

We define matrices $\mathbb{O}$ with $O_{jk} = \delta_{jk} \Omega_{kH}$ and $\tilde{\mathbb{S}}$ with $\tilde{S}_{jk} = \int d\mathbf{r} \psi_j^* \tilde{S} \psi_k$. Thus $d_{Nj} = [\mathbb{D}_N]_j$ where $\mathbb{D}_N = (2\mathbb{O} + \tilde{\mathbb{S}}) \mathbb{P}_N$ and $\mathbb{P}_N$ is a solution of (91). We can use this expansion to

calculate the Coulomb projections W_{NM} defined in (37). Using (100) we find

$$\begin{aligned} W_{NM} &= \mathbb{D}_M^\dagger \mathbb{W}^{(\psi)} \mathbb{D}_N \\ &= \mathbb{P}_M^\dagger (2\mathbb{O} + \tilde{\mathbb{S}}) \mathbb{W}^{(\psi)} (2\mathbb{O} + \tilde{\mathbb{S}}) \mathbb{P}_N \end{aligned} \quad (101)$$

where

$$[\mathbb{W}^{(\psi)}]_{jk} = \int d\mathbf{r} d\mathbf{r}' \frac{\Psi^0(\mathbf{r}) \psi_j^*(\mathbf{r}) \Psi^0(\mathbf{r}') \psi_k(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}. \quad (102)$$

We can use W_{NM} to calculate the correlation energy through equation (38).

In molecular orbital language it is clear the calculation of $\mathbb{W}^{(\psi)}$ is the only step involving four-center integrals required by a correlation energy calculation. However, in contrast to the four orbital (two occupied, two unoccupied) terms ($ia|jb$) required by a full KS calculation it requires two uncontracted orbital indices (occupied or unoccupied) only. For large molecules this is a substantial saving.

IX. 1D HARMONIC OSCILLATOR

We define a 1D non-interacting and spin-free Harmanoic oscillator with N_e electrons in the groundstate and $V^{\text{KS}}(x) = x^2/2$. Here the KS orbital wavefunctions and energies take the form

$$\psi_j(x) = \kappa_j H_j(x) e^{-x^2/2}, \quad \epsilon_j = \frac{2j+1}{2} \quad (103)$$

where $H_j(x)$ are Hermite polynomials with $j \geq 0$ and $\kappa_j = [\sqrt{\pi} 2^j j!]^{-1/2}$. Since the orbitals are filled up to $j \leq N_e - 1$ the density is

$$n^0(x) = \sum_{j=0}^{N_e-1} \frac{H_j(x)^2 e^{-x^2}}{\sqrt{\pi} 2^j j!} \quad (104)$$

and it is clear from the expansion of H_{N_e-1} that $\lim_{x \rightarrow \infty} n^0(x) \sim x^{2(N_e-1)} e^{-x^2}$ and $\lim_{x \rightarrow \infty} \eta(x) \sim -x$. Thus the system is well-bounded and the KS-orbital approach should work well.

In the one-electron case $\eta = -x$, $t = \frac{1}{2}$ and $V_{,xx} = 1$. and we can solve everything analytically. The matrices required to form $\mathbb{N}$ and $\mathbb{R}$ in equations (92) and (93) have elements

$T_{jk} = \frac{1}{2}\delta_{jk}$, $V_{jk} = \delta_{jk}$ and

$$\begin{aligned} D_{jk} &= \kappa_j \kappa_k \int dx e^{-x^2/2} H_j(x) [(\partial_x + x) H_k(x) e^{-x^2/2}] \\ &= 2k \kappa_j \kappa_k \int dx e^{-x^2} H_j(x) H_{k-1}(x) \\ &= \sqrt{2(j+1)} \delta_{j(k-1)}. \end{aligned}$$

Here $[\mathbb{D}^\dagger \mathbb{D}]_{jk} = 2j\delta_{jk}$, $[\mathbb{D}^{\dagger 2} \mathbb{D}^2]_{jk} = 4j(j-1)\delta_{jk}$ and $[\mathbb{D}^{\dagger 3} \mathbb{D}^3]_{jk} = 8j(j-1)(j-2)\delta_{jk}$.

Thus $N_{jk} = 2j\delta_{jk}$ and $R_{jk} = 2j\delta_{jk} + 6j(j-1)\delta_{jk} + 2j(j-1)(j-2)\delta_{jk} = 2j^3\delta_{jk}$. We can ignore the $j = 0$ solution as it will not contribute to $\hat{\chi}_0$. We thus choose solutions with $N_j > 0$ where $p_{N_j j} = \delta_{N_j j} / \sqrt{2j}$. Here $\Omega_{N_j} = \frac{2j+1}{2} - \frac{1}{2} = j$ as expected and the transition density mode is $d_{N_j}(x) = -\psi_0(x) [(\partial_x^2 + 1 - x^2)\psi_j(x)] / \sqrt{2j} = \sqrt{2\Omega_{N_j}} \psi_0(x) \psi_j(x)$.

In the many-electron case we can solve the problem semi-analytically for much of it, requiring numerics only for terms involving $1/n^0$ and subsequent diagonalisations. Errors occur due to truncation of the basis set but integrals can be obtained with near exact accuracy with Gauss-Hermite quadrature. It should be noted that the two-electron system is not predicted exactly by the CM as there is no spin degeneracy.

The rapid convergence of the method is demonstrated in Table I where we show the convergence of the fourth transition frequency Ω_4 for the two-electron system. With as few as 15 states the error is already under one part in ten thousand.

N_{Bas}	5	10	15	20	50	∞
Ω_4	4.1105	3.8797	3.8805	3.8801	3.8802	3.8802
$\log_{10} \text{Err} $	-1.22	-3.95	-4.05	-5.06	-7.49	$-\infty$

TABLE I. Convergence of the fourth transition frequency for the two electron system. Here $\text{Err} = \Omega_4^{N_{\text{Bas}}} / \Omega_4^\infty - 1$.

The KS transition frequencies of an N_e electron system have energies $\Omega_{ai} = \frac{2a+1}{2} - \frac{2i+1}{2} = J$ and each frequency J has multiple contributing transitions. Here $\min(J, N_e)$ modes have transition frequency J and the density transitions are proportional to $\psi_i(x)\psi_{i+J}(x)$ when $i \leq N_e$ and $i + J > N_e$.

We present the CM transition frequencies in Table II for systems with up to 20 electrons. The CM has *single-valued* frequencies distributed approximately with the integers. Each CM transition density is therefore composed of multiple KS-like transitions which, by the

sum rules (32) must have their weights h_{aiN} dominated by transitions with $\Omega_{ai} \approx \Omega_N$. In Table III we show some weights h_{aiN} for the five electron system. It is clear that the N th CM transition puts most weight on the HOMO-HOMO+ N [$5 - (5 + N)$] KS transition as one would hope.

N	Transition frequencies Ω_N				
	$N_e = 1$	$N_e = 2$	$N_e = 5$	$N_e = 10$	$N_e = 20$
1	1.0000	1.0000	1.0000	1.0000	1.0000
2	2.0000	2.0000	2.0000	2.0000	2.0000
3	3.0000	3.0000	3.0000	3.0000	3.0000
4	4.0000	3.8802	3.9531	3.9859	3.9963
5	5.0000	4.8680	4.8162	4.9225	4.9772
6	6.0000	5.7877	5.6869	5.7886	5.9216
7	7.0000	6.7689	6.6309	6.6381	6.8128
8	8.0000	7.7154	7.5079	7.5381	7.6683
10	10.0000	9.6579	9.3578	9.3030	9.4403
20	20.0000	19.4882	18.6483	18.2017	18.1953

TABLE II. First eight, 10th and 20th distinct eigenfrequencies of the KS system and CM system with different numbers of electrons N_e . The KS transitions are all integer and degenerate with $\min(\Omega, N_e)$ transitions for a given integer frequency.

It is remarkable that the lowest three frequencies are exact to four decimal places for $N_e > 1$, particularly as the second smallest is comprised of two transitions and the third of three in all but the two-electron case. By the sum rules (32) this means that each mode must be comprised solely of transitions with the given frequency, and as a consequence, both Ω_N and d_N are predicted exactly for the HOMO-LUMO transition. That such a relationship holds for as many as 20 electrons demonstrates a strength of the physics and a resilience of the approximations in the CM approach.

X. CONCLUSION

In this paper we have reformulated, simplified and investigated the CM formalism of refs 4–7, and provided a direct proof of its exactness in one-electron systems. We have described two formulations: a normalised approach very similar to the usual form, and a scalar approximation that is directly equivalent to the usual tensor form for one-electron and one-dimensional systems.

The orthonormal form described in Section VI is vital for the description and calculation of localised systems, where the standard $\mathbf{u}$ -based formulation does not behave well. This reformulation is required for calculation of atomic, and molecular systems using standard basis set approaches involving GTO and STOs. It also provides a relatively simple way of proving (see Section VII), direct from the Schrödinger equation, that the CM is exact for

N	Ω_N	(i,a)	$h_{aiN}\%$	(i,a)	$h_{aiN}\%$	(i,a)	$h_{aiN}\%$
1	1.0000	(5,6)	100.0				
2	2.0000	(5,7)	60.0	(4,6)	40.0		
3	3.0000	(5,8)	53.8	(4,7)	30.8	(3,6)	15.4
4	3.9531	(5,9)	54.4	(4,8)	27.1	(3,7)	11.5
		(2,6)	3.8	(4,6)	1.9	(5,7)	1.2
5	4.8162	(5,10)	52.8	(4,9)	23.9	(3,8)	9.0
		(3,6)	6.2	(5,8)	4.2	(2,7)	2.6
7	6.6309	(5,12)	49.8	(4,11)	20.0	(5,10)	6.8
		(3,10)	6.7	(3,8)	5.7	(2,7)	5.2
		(1,6)	1.9	(2,9)	1.7		
10	9.3578	(5,15)	45.0	(4,14)	16.5	(5,13)	14.2
		(3,13)	5.0	(3,11)	4.0	(2,10)	3.2
		(2,8)	3.1	(1,7)	2.3	(5,17)	1.1
		(2,12)	1.1				

TABLE III. Weights h_{aiN} in % with $> 1\%$ contribution for the five electron system. The KS transition frequencies are $\Omega_{ai} = a - i$. Tabulated weights may not sum to 100% due to the absence of $< 1\%$ contributions.

one-electron systems.

The scalar approximation of Section VIII provides a simplified approach to the CM, with some sacrifice of some exact properties of the full theory in general systems. It could be used as a faster alternative itself or as a doorway to further and cruder approximations. In particular it opens up analysis of transition frequencies and densities in a simple manner by relating information from the KS orbitals to the collective modes in the CM. The approximation is exact in one-electron systems and should accurately predict the physics of asymptotic regions where the behaviour is essentially one-electron like.

In this approximation, at least for the difficult four-center term required by correlation energy calculations, the scalar CM approximation has an efficiency gain of $O(N_e^2)$ over regular KS where N_e is the number of occupied electronic orbitals. This means that in large molecules (and in periodic bulks), evaluation of E_c under the approximate CM will be a significantly quicker calculation, with corresponding lower memory requirements. The approximation also makes a gain of $O(3^3)$ in diagonalisation compared to the full CM theory and $O(3^2)$ in the four-center terms.

Both reformulations presented here will aid in the development of a basis set based, CM approach for atomic and molecular systems. The difficulty in such an approach is the presence of non-constant denominators in $t_{\mu\nu}$, η_α and S . We believe, however, that a tractable approach to dealing with these should be possible by using Gauss-Hermite quadrature with GTOs.

It should also be possible to further develop vdW functionals that use a further approximation to the CM based on cleverly chosen orbital-like basis functions, possibly incorporating some unoccupied KS orbitals. Such functionals could incorporate ideas from refs 15–17 to be made even more efficient than the functional described here or in ref. 7, without sacrificing the vital physics that make the CM attractive.

ACKNOWLEDGMENTS

We would like to thank Janos Angyan, Andreas Savin and Giovanni Vignale for much fruitful discussion. G. J. thanks John Dobson for his hospitality and support during a sabbatical stay in Brisbane. J.F.D. and T.G. were supported by ARC Discovery Grant DP1096240. I. V. T. was supported by the Spanish MICINN, Grant No. FIS2010-21282-

Appendix A: Kinetic stress tensor

There is some debate over the appropriate form of the kinetic stress tensor $\mathbf{T}$. We have chosen the form given in (3) and (4), whereas the original CM papers used the form given in (13). One requirement is that, for Kohn-Sham systems, $\mathbf{T}^0$ obeys the force balance condition $\nabla \cdot \mathbf{T}^0 = -n^0 \nabla V^{\text{KS}}$ which is true both for the form (4) used in this manuscript and for equation 17 of ref. 5 [leading to our equation (14) for a Kohn-Sham groundstate].

There are various ways to arrive at the possible forms of $\mathbf{T}(\mathbf{r})$ (see e.g. refs 29–33). For completeness, we show one way to arrive at the form (3), (4) chosen for the present work. This form helps to make the CM equations particularly simple. We base our form on the Wigner transform of the “classical” kinetic stress tensor. In a “classical” fluid with no current present, $\mathbf{T}$ is defined by

$$T_{\mu\nu}(\mathbf{r}) = \int d\mathbf{p} p_\mu p_\nu f(\mathbf{r}, \mathbf{p}). \quad (\text{A1})$$

Here f is the classical one-body distribution function - i.e. the phase space probability density for finding a particle at position $\mathbf{r}$ with momentum $\mathbf{p}$. To model a quantum system we can replace $f(\mathbf{r}, \mathbf{p})$ by the Wigner form

$$f(\mathbf{r}, \mathbf{p}) = \int \frac{d\mathbf{x}}{(2\pi)^3} \rho\left(\mathbf{r} + \frac{\mathbf{x}}{2}, \mathbf{r} - \frac{\mathbf{x}}{2}\right) e^{i\mathbf{p} \cdot \mathbf{x}} \quad (\text{A2})$$

where $\rho(\mathbf{r}, \mathbf{r}')$ is the quantal one-body density matrix. Using the short-hand $\rho = \rho(\mathbf{r} + \mathbf{x}/2, \mathbf{r} - \mathbf{x}/2)$ we thus find

$$\begin{aligned} T_{\mu\nu}(\mathbf{r}) &= \int d\mathbf{p} \int \frac{d\mathbf{x}}{(2\pi)^3} \rho p_\mu p_\nu e^{i\mathbf{p} \cdot \mathbf{x}} \\ &= \int d\mathbf{p} \int \frac{d\mathbf{x}}{(2\pi)^3} \rho [-\partial_{x_\mu} \partial_{x_\nu} e^{i\mathbf{p} \cdot \mathbf{x}}] \\ &= \int \frac{d\mathbf{x}}{(2\pi)^3} [-\partial_{x_\mu} \partial_{x_\nu} \rho] \int d\mathbf{p} e^{i\mathbf{p} \cdot \mathbf{x}} \\ &= [-\partial_{x_\mu} \partial_{x_\nu} \rho]_{\mathbf{x}=\mathbf{0}}, \end{aligned} \quad (\text{A3})$$

and so, by the chain rule,

$$\begin{aligned} T_{\mu\nu}(\mathbf{r}) &= -\partial_{x_\mu} \partial_{x_\nu} \rho(\mathbf{r} + \mathbf{x}/2, \mathbf{r} - \mathbf{x}/2)|_{\mathbf{x}=\mathbf{0}} \\ &= \frac{1}{4} [(\partial_\mu \partial'_\nu + \partial'_\mu \partial_\nu - \partial_\mu \partial_\nu - \partial'_\mu \partial'_\nu) \rho(\mathbf{r}, \mathbf{r}')]_{\mathbf{r}=\mathbf{r}'} \end{aligned} \quad (\text{A4})$$

with $\partial_\mu \equiv \partial_{r_\mu}$ and $\partial'_\mu \equiv \partial_{r'_\mu}$. Furthermore

$$\begin{aligned} & \frac{1}{4} [(\partial_\mu \partial_\nu + \partial_\mu \partial'_\nu + \partial'_\mu \partial_\nu + \partial'_\mu \partial'_\nu) \rho(\mathbf{r}, \mathbf{r}')]_{\mathbf{r}=\mathbf{r}'} \\ &= \frac{1}{4} \partial_\mu \partial_\nu \rho(\mathbf{r}, \mathbf{r}) = \frac{1}{4} \partial_\mu \partial_\nu n(\mathbf{r}) \end{aligned} \quad (\text{A5})$$

Combining (A4) and (A5) we find

$$\begin{aligned} T_{\mu\nu}(\mathbf{r}) &= \frac{1}{2} [(\partial_\mu \partial'_\nu + \partial'_\mu \partial_\nu) \rho(\mathbf{r}, \mathbf{r}')]_{\mathbf{r}=\mathbf{r}'} \\ &\quad - \frac{1}{4} \partial_\mu \partial_\nu n(\mathbf{r}). \end{aligned} \quad (\text{A6})$$

for a state with zero current, in agreement with (3).

Since $\rho(\mathbf{r}, \mathbf{r}') = \sum_i f_i \psi_i^*(\mathbf{r}) \psi_i(\mathbf{r}')$ in a groundstate KS system we find the stress tensor T^0 for this situation to be

$$\begin{aligned} T_{\mu\nu}^0(\mathbf{r}) &= \frac{1}{2} \sum_i f_i (\partial_\mu \partial'_\nu + \partial'_\mu \partial_\nu) \psi_i^*(\mathbf{r}) \psi_i(\mathbf{r}')|_{\mathbf{r}=\mathbf{r}'} \\ &\quad - \frac{1}{4} [\partial_\mu \partial_\nu n^0(\mathbf{r})] \\ &= \Re \sum_i f_i [\partial_\mu \psi_i^*] [\partial_\nu \psi_i] - \frac{1}{4} [\partial_\mu \partial_\nu n^0] \end{aligned} \quad (\text{A7})$$

Thus (3) is exactly the Wigner version of the “classical” kinetic stress tensor (A1).

Using (5) we demonstrate that $\partial_\alpha T_{\alpha\mu}^0 = -n^0 V_{,\mu}^{\text{KS}}$. Here we note that

$$\partial_\alpha T_{\alpha\mu}^0 = \frac{1}{2} \sum_j f_j t_{j\mu} \quad (\text{A8})$$

where

$$\begin{aligned} t_{j\mu} &= \Re \{ [\nabla^2 \psi_j^*] [\partial_\mu \psi_j] + [\partial_\alpha \psi_j^*] [\partial_\alpha \partial_\mu \psi_j] \\ &\quad - [\partial_\alpha \psi_j^*] [\partial_\alpha \partial_\mu \psi_j] - \psi_j^* [\partial_\mu \nabla^2 \psi_j] \} \\ &= \Re \{ [\nabla^2 \psi_j^*] [\partial_\mu \psi_j] - \psi_j^* [\partial_\mu \nabla^2 \psi_j] \} \\ &= -2V_{,\mu}^{\text{KS}} |\psi_j|^2 \end{aligned}$$

and we have used the Schrödinger equation $\nabla^2 \psi_j = 2(V^{\text{KS}} - \epsilon_j) \psi_j$ to derive the final expression. Finally $\partial_\alpha T_{\mu\alpha}^0 = -\sum_j f_j V_{,\mu}^{\text{KS}} |\psi_j|^2 = -n^0 V_{,\mu}^{\text{KS}}$ and the proof is complete.

Comparing this with the form $\bar{T}^0$ (14) used in the original CM formulation, we find $\partial_\alpha \bar{T}_{\mu\alpha}^0 = \partial_\alpha T_{\mu\alpha}^0$ since $\partial_\alpha [\partial_\mu \partial_\alpha n^0] = [\partial_\mu \nabla^2 n^0] = \partial_\alpha \delta_{\mu\alpha} [\nabla^2 n^0]$. Since these are precisely the terms that differ between (4) and (14) it follows that the gradients must be identical.

Appendix B: Compact form of the CM equations

In an earlier work⁷ we state without proof that equation (8) defined via (11) here (equations 2-5 of ref. 7) are equivalent to equations 14-16 of ref. 5. The only non-notational difference is in the kinetic force term $\mathbf{F}^{\text{1Kin}} = -\hat{K}\mathbf{u}$ defined here via (9), in equation 14 of ref. 5, and in equation 53 of ref. 6 (abbreviated as G53) where it is derived from their equation C8 (abbreviated as GC8).

We demonstrate that the two forms are equivalent by working from GC8. While the same result can be obtained directly from G53 the derivation is less clear and less succinct. Specifically we must show that

$$\mathbf{F}_\mu^{\text{1Kin}} \equiv \frac{\delta \mathcal{T}_2[\mathbf{u}]}{\delta u_\mu} = -\hat{K}_{\mu\nu} u_\nu \quad (\text{B1})$$

where $\hat{K}_{\mu\nu}$ is defined in (9), since the remaining terms in (8) follow directly from equations 14-16 of ref. 5.

Following Gao et al⁶ we write GC8 as $\mathcal{T}_2[\mathbf{u}] = \int d\mathbf{r} \mathcal{I}$ where

$$\begin{aligned} \mathcal{I} = & \left\{ \mathcal{K}_{\mu\nu} (4U_{\mu\alpha} U_{\nu\alpha} - u_{\alpha,\mu} u_{\alpha,\nu}) + \frac{n^0}{8} U_{\alpha\alpha,\mu} U_{\nu\nu,\mu} \right. \\ & \left. + \frac{n^0_{,\nu}}{2} U_{\mu\nu} \partial_\mu U_{\alpha\alpha} + \frac{n^0_{,\nu}}{4} u_{\alpha,\mu} \partial_\nu u_{\mu,\alpha} \right\}, \end{aligned} \quad (\text{B2})$$

with $U_{\mu\nu} = \frac{1}{2}(u_{\mu,\nu} + u_{\nu,\mu})$, $U_{\alpha\alpha} = u_{\alpha,\alpha}$, and $\mathcal{K}_{\mu\nu} = \frac{1}{2}\Re \sum_i f_i [\partial_\mu \psi_i^*] [\partial_\nu \psi_i]$.

We can use integration by parts to remove derivatives of n^0 from the integrand $\mathcal{I}$. As such

$$\begin{aligned} \mathcal{T}_2[\mathbf{u}] = & \frac{1}{2} \int d\mathbf{r} \left(2\mathcal{K}_{\mu\nu} \mathcal{Y}_{\mu\nu} - \frac{n^0}{4} \mathcal{Z} \right) \\ = & \frac{1}{2} \int d\mathbf{r} \left[T_{\mu\nu}^0 \mathcal{Y}_{\mu\nu} + \frac{n^0}{4} (\mathcal{Y}_{\mu\nu,\mu\nu} - \mathcal{Z}) \right] \end{aligned} \quad (\text{B3})$$

where we define $T_{\mu\nu}^0 = 2\mathcal{K}_{\mu\nu} - \frac{1}{4}n^0_{,\mu\nu}$ to be equal to (4) rather than the form appearing in refs 5 and 6. The undefined terms in the integrand of (B3) take the form

$$\mathcal{Y}_{\mu\nu} = u_{\mu,\alpha} u_{\nu,\alpha} + u_{\alpha,\mu} u_{\nu,\alpha} + u_{\mu,\alpha} u_{\alpha,\nu}, \quad (\text{B4})$$

$$\begin{aligned} \mathcal{Z} = & u_{\mu,\mu\alpha} u_{\nu,\nu\alpha} + 2u_{\mu,\nu\alpha} u_{\nu,\mu\alpha} \\ & + u_{\mu,\nu\alpha\alpha} u_{\nu,\mu} + u_{\mu,\nu} u_{\nu,\mu\alpha\alpha} \\ & + u_{\mu,\alpha\alpha} u_{\nu,\mu\nu} + u_{\mu,\nu\mu} u_{\nu,\alpha\alpha} \\ & + 2u_{\mu,\mu\nu\alpha} u_{\nu,\alpha} + 2u_{\mu,\alpha} u_{\nu,\mu\nu\alpha} \end{aligned} \quad (\text{B5})$$

where we have expanded $U_{\mu\nu}$ and used the product rule on all derivatives to arrive at these forms. The following identities are also used in the derivation of $\mathcal{Z}$:

$$\begin{aligned}
U_{\alpha\alpha,\mu}U_{\nu\nu,\mu} &= u_{\mu,\mu\alpha}u_{\nu,\nu\alpha} \\
2\partial_\nu u_{\mu,\nu}\partial_\mu u_{\alpha,\alpha} &= \partial_\alpha u_{\mu,\alpha}\partial_\mu u_{\nu,\nu} + \partial_\alpha u_{\nu,\alpha}\partial_\nu u_{\mu,\mu}, \\
2\partial_\nu u_{\nu,\mu}\partial_\mu u_{\alpha,\alpha} &= \partial_\nu u_{\nu,\alpha}\partial_\alpha u_{\mu,\mu} + \partial_\mu u_{\mu,\alpha}\partial_\alpha u_{\nu,\nu}, \\
2\partial_\nu u_{\alpha,\mu}\partial_\nu u_{\mu,\alpha} &= \partial_\alpha u_{\nu,\mu}\partial_\alpha u_{\mu,\nu} + \partial_\alpha u_{\mu,\nu}\partial_\alpha u_{\nu,\mu}
\end{aligned}$$

which follow from exchange of indices under summation (eg. $A_{\nu\mu}B_{\mu\nu} \equiv A_{\mu\nu}B_{\nu\mu}$).

We can expand the terms of $\mathcal{Y}_{\mu\nu,\mu\nu}$ as follows:

$$\begin{aligned}
\partial_\mu\partial_\nu u_{\mu,\alpha}u_{\nu,\alpha} &= u_{\mu,\mu\alpha}u_{\nu,\nu\alpha} + u_{\mu,\nu\alpha}u_{\nu,\mu\alpha} \\
&\quad + u_{\mu,\mu\nu\alpha}u_{\nu,\alpha} + u_{\mu,\alpha}u_{\nu,\mu\nu\alpha}, \\
\partial_\mu\partial_\nu u_{\mu,\alpha}u_{\alpha,\nu} &= u_{\mu,\mu\nu}u_{\nu,\alpha\alpha} + u_{\mu,\nu\alpha}u_{\nu,\mu\alpha} \\
&\quad + u_{\mu,\mu\nu\alpha}u_{\nu,\alpha} + u_{\mu,\nu}u_{\nu,\mu\alpha\alpha}, \\
\partial_\mu\partial_\nu u_{\nu,\alpha}u_{\alpha,\mu} &= u_{\mu,\alpha\alpha}u_{\nu,\mu\nu} + u_{\mu,\nu\alpha}u_{\nu,\mu\alpha} \\
&\quad + u_{\mu,\alpha}u_{\nu,\mu\nu\alpha} + u_{\mu,\nu\alpha\alpha}u_{\nu,\mu}
\end{aligned}$$

where we again exchange indices where appropriate. This leads to the following result

$$\begin{aligned}
\mathcal{Y}_{\mu\nu,\mu\nu} &= u_{\mu,\mu\alpha}u_{\nu,\nu\alpha} + 3u_{\mu,\nu\alpha}u_{\nu,\mu\alpha} \\
&\quad + u_{\mu,\alpha\alpha}u_{\nu,\mu\nu} + u_{\mu,\mu\nu}u_{\nu,\alpha\alpha} \\
&\quad + u_{\mu,\nu\alpha\alpha}u_{\nu,\mu} + u_{\mu,\nu}u_{\nu,\mu\alpha\alpha} \\
&\quad + 2u_{\mu,\alpha}u_{\nu,\mu\nu\alpha} + 2u_{\mu,\mu\nu\alpha}u_{\nu,\alpha}
\end{aligned} \tag{B6}$$

$$= \mathcal{Z} + u_{\mu,\nu\alpha}u_{\nu,\mu\alpha} \tag{B7}$$

and thus $\mathcal{Y}_{\mu\nu,\mu\nu} - \mathcal{Z} = u_{\nu,\mu\alpha}u_{\mu,\nu\alpha}$. The cancellation of so many terms is quite remarkable.

Finally (B3) becomes

$$\begin{aligned}
\mathcal{T}_2[\mathbf{u}] &= \frac{1}{2} \int d\mathbf{r} \left\{ T_{\mu\nu}^0 (u_{\mu,\alpha}u_{\nu,\alpha} + u_{\alpha,\mu}u_{\nu,\alpha} + u_{\mu,\alpha}u_{\alpha,\nu}) \right. \\
&\quad \left. + (n^0/4)u_{\nu,\mu\alpha}u_{\mu,\nu\alpha} \right\}
\end{aligned} \tag{B8}$$

$$= -\frac{1}{2} \int d\mathbf{r} u_\mu \hat{K}_{\mu\nu} u_\nu \tag{B9}$$

$$\begin{aligned}
\hat{K}_{\mu\nu} &= -\frac{1}{4} \partial_\nu \partial_\alpha n^0 \partial_\alpha \partial_\mu \\
&\quad + \partial_\alpha T_{\mu\nu}^0 \partial_\alpha + \partial_\alpha T_{\alpha\nu}^0 \partial_\mu + \partial_\nu T_{\alpha\mu}^0 \partial_\alpha.
\end{aligned} \tag{B10}$$

Here we used integration by parts on the derivatives of u_μ to obtain (B9) and (B10). The operator $\hat{K}$ defined in (B10) is identical to that defined in (9). Taking the functional derivative w.r.t. u_μ thus gives

$$\mathbf{F}_\mu^{1\text{Kin}} = \frac{\delta \mathcal{T}_2[\mathbf{u}]}{\delta u_\mu} = -\hat{K}_{\mu\nu} u_\nu \quad (\text{B11})$$

and it is clear that (B1) is satisfied.

Appendix C: Correlation energy expressions

Let us first define the projected Coulomb operator $W_{NM} = \langle \mathbf{u}_M | \hat{Q} | \mathbf{u}_N \rangle$ where $\hat{Q}$ is defined in (16) or (17). This can be written as

$$W_{NM} = \int d\mathbf{r} \mathbf{u}_M^*(\mathbf{r}) \cdot [\hat{Q} \mathbf{u}_N](\mathbf{r}) \quad (\text{C1})$$

$$= \int d\mathbf{r} d\mathbf{r}' \mathbf{u}_M^*(\mathbf{r}) \cdot \mathbf{Q}(\mathbf{r}, \mathbf{r}') \cdot \mathbf{u}_N(\mathbf{r}') \quad (\text{C2})$$

$$= \int \frac{d\mathbf{r} d\mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|} d_M^*(\mathbf{r}) d_N(\mathbf{r}'). \quad (\text{C3})$$

where similar equivalences hold true for $W_{NM\lambda} = \langle \mathbf{u}_{M\lambda} | \hat{Q} | \mathbf{u}_{N\lambda} \rangle$ or for alternative forms of the Coulomb potential (e.g. range-separated).

Using equations (18)-(19) and (24)-(25) we can write the correlation energy as

$$E_c = \frac{1}{2} \int_0^1 d\lambda \int_0^\infty \frac{d\sigma}{\pi} \int \frac{d\mathbf{r} d\mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|} \times \sum_N \left(\frac{d_{N\lambda}^*(\mathbf{r}) d_{N\lambda}(\mathbf{r}')}{\Omega_{N\lambda}^2 + \sigma^2} - \frac{d_N^*(\mathbf{r}) d_N(\mathbf{r}')}{\Omega_N^2 + \sigma^2} \right) \quad (\text{C4})$$

$$= \frac{1}{2} \int_0^1 d\lambda \int_0^\infty \frac{d\sigma}{\pi} \sum_N \left(\frac{W_{NN\lambda}}{\Omega_{N\lambda}^2 + \sigma^2} - \frac{W_{NN}}{\Omega_N^2 + \sigma^2} \right) \quad (\text{C5})$$

$$= \frac{1}{2} \int_0^1 d\lambda \sum_N \left(\frac{W_{NN\lambda}}{2\Omega_{N\lambda}} - \frac{W_{NN}}{2\Omega_N} \right). \quad (\text{C6})$$

Here the governing eigen-equations for $\mathbf{u}_{N\lambda}$ are as defined in (19)

$$\Omega_{N\lambda}^2 n^0 \mathbf{u}_{N\lambda} = (\hat{R} + \lambda \hat{Q}) \mathbf{u}_{N\lambda} \quad (\text{C7})$$

with normalisation $\int n^0(\mathbf{r}) \mathbf{u}_{N\lambda}^*(\mathbf{r}) \cdot \mathbf{u}_{M\lambda}(\mathbf{r}) = \delta_{NM}$.

Following the ideas of Furche,²⁵ we can take the λ derivative of $\int d\mathbf{r} \mathbf{u}_{N\lambda} \cdot (\text{C7})$ to work directly from (C6). Here

$$2\Omega_{N\lambda} [\partial_\lambda \Omega_{N\lambda}] = \int \mathbf{u}_{N\lambda}^* \cdot \hat{Q} \mathbf{u}_{N\lambda}. \quad (\text{C8})$$

where derivatives of $\mathbf{u}_{N\lambda}$ can be ignored by the Hellman-Feynman theorem. Thus

$$[\partial_\lambda \Omega_{N\lambda}] = \int d\mathbf{r} \frac{\mathbf{u}_{N\lambda}^*(\mathbf{r})[\hat{Q}\mathbf{u}_{N\lambda}](\mathbf{r})}{2\Omega_{N\lambda}} = \frac{W_{NN\lambda}}{2\Omega_{N\lambda}}. \quad (\text{C9})$$

and we can write the correlation as a sum over zero-point energies such that

$$E_c = \frac{1}{2} \int_0^1 d\lambda \sum_N \left([\partial_\lambda \Omega_{N\lambda}] - \frac{W_{NN}}{2\Omega_N} \right) \quad (\text{C10})$$

$$= \frac{1}{2} \sum_N \left[\bar{\Omega}_N - \Omega_N \left(1 + \frac{W_{NN}}{2\Omega_N^2} \right) \right] \quad (\text{C11})$$

where $\bar{\Omega}_N = \Omega_{N1}$. In certain systems it may make sense to work with exchange and correlation together. Here

$$E_{xc} = \frac{1}{2} \sum_N [\bar{\Omega}_N - \Omega_N] - \frac{1}{2} \int d\mathbf{r} n^0(\mathbf{r}) w_C(\mathbf{r}) \quad (\text{C12})$$

where $w_C(\mathbf{r}) = \int d\mathbf{r}' \frac{\delta(\mathbf{r}-\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|}$ is like the Coulomb potential at zero distance.

To solve for $\bar{\Omega}_N$ involves a difficult diagonalisation of $\hat{\mathbf{R}} + \hat{\mathbf{Q}}$ and may be best avoided. We can use the complete and orthogonal nature of $\mathbf{u}_{N\lambda}$ and $\mathbf{u}_N$ to write $\mathbf{u}_{N\lambda} = \sum_K U_{NK} \mathbf{u}_K$ where $\mathbb{U}^\dagger \mathbb{U} = \mathbb{I}$. Thus we can write

$$\chi_\lambda(\mathbf{r}, \mathbf{r}') = -\Re \sum_{NM} X_{NM\lambda} d_N^*(\mathbf{r}) d_M(\mathbf{r}') \quad (\text{C13})$$

where $X_{NM0} = \delta_{NM}/(\sigma^2 + \Omega_N^2)$ or $\mathbb{X}_0 = (\sigma^2 + \mathbb{L})^{-1}$ where $L_{NM} = \delta_{NM} \Omega_N^2$. Solving for $\hat{\chi}_\lambda = \hat{\chi}_0 + \lambda \hat{\chi}_0 \hat{v} \hat{\chi}_\lambda$ gives

$$\mathbb{X}_\lambda = \mathbb{X}_0 - \lambda \mathbb{X}_0 \mathbb{W} \mathbb{X}_\lambda, \quad \mathbb{X}_\lambda = \frac{\mathbb{I}}{\sigma^2 + \mathbb{L} + \lambda \mathbb{W}}. \quad (\text{C14})$$

This corresponds to

$$E_c = \frac{1}{2} \int_0^1 d\lambda \int_0^\infty \frac{d\sigma}{\pi} \times \text{Tr} \left[\frac{\mathbb{W}}{\sigma^2 + \mathbb{L} + \lambda \mathbb{W}} - \frac{\mathbb{W}}{\sigma^2 + \mathbb{L}} \right] \quad (\text{C15})$$

which can sometimes prove useful in real calculations

We can relate all this back to (C11). Solving the eigen-equation $\mathbb{V} \mathbb{D}_\lambda = (\mathbb{L} + \lambda \mathbb{W}) \mathbb{V}$ gives

$$X_{MN\lambda} = [\mathbb{V}(\sigma^2 + \mathbb{D}_\lambda)^{-1} \mathbb{V}^\dagger]_{MN} \quad (\text{C16})$$

$$= \sum_K U_{KM}^* (\sigma^2 + \Omega_{K\lambda}^2)^{-1} U_{KN}. \quad (\text{C17})$$

With appropriate sorting of the eigen-value/vector pairs it is clear that the eigenvalues D_{NN1} of $\mathbb{L} + \mathbb{W}$ are $\bar{\Omega}_N^2 = \Omega_{N1}^2$ and $\mathbb{U} = \mathbb{V}^\dagger$. Thus the two expressions are equivalent.

Furthermore for $\Omega_N^2 \gg \bar{W}_N$ where $\bar{W}_N = \sum_M |W_{NM}|$ we can solve the eigen-problem perturbatively so that $\bar{\Omega}_N^2 \approx \Omega_N^2 + W_{NN}$. Thus $\bar{\Omega}_N = \sqrt{\Omega_N^2 + W_{NN}} \approx \Omega_N(1 + \frac{W_{NN}}{2\Omega_N^2} - \frac{W_{NN}^2}{8\Omega_N^4})$. We can speed calculation and improve numerical stability by choosing an N^* above which we use the approximation. Setting $\beta_N = \frac{W_{NN}}{2\Omega_N^2}$ we then find

$$E_c \approx \frac{1}{2} \sum_{N=0}^{N^*} [\bar{\Omega}_N - \Omega_N(1 + \beta_N)] - \sum_{N > N^*} \frac{\Omega_N \beta_N^2}{4}. \quad (\text{C18})$$

Such a perturbative approach will be almost guaranteed convergent if we replace the Coulomb potential by the range-separated³⁴⁻³⁶ form $V_C^{(qrs)}(R) = \text{erf}(q_{RS}R)/R$.

Appendix D: One-Electron governing operator

As discussed in section VII [equations (61)-(67)] we can obtain the exact linear perturbation solutions in a one-electron system via a solution of

$$\omega^2 \xi = \hat{R}^{1e^-} \xi \quad (\text{D1})$$

where the associated change in density is $n^1 = -\nabla \psi \xi^{1e^-}$. Since this is an identical problem to finding the CM solutions via

$$\omega^2 \xi = \hat{R}^{(\xi)} \xi \quad (\text{D2})$$

it is clear that if $\hat{R}^{1e^-} = \hat{R}^{(\xi)}$ then the CM is exact for one-electron systems.

For a one-electron system we set $V = V^{\text{KS}} - \epsilon_0$ such that the groundstate occupied orbital ψ is a solution of $[-\frac{1}{2}\nabla^2 + V]\psi = 0$. Equations (55) thus become $\eta_\alpha = \partial_\alpha \log |\psi|$ and $S = 2V$ and thus $\hat{h} = -\frac{1}{2}(\nabla^2 - S)$. The former can be used to derive the following

$$\begin{aligned} [\partial_\mu \eta_\nu] &= [\partial_\nu \eta_\mu] = -2t_{\mu\nu} \\ [\nabla^2 \eta_\mu] &= -2[\partial_\alpha t_{\alpha\mu}] = S_{,\mu} + 4\eta_\alpha t_{\alpha\mu} \end{aligned}$$

where these identities will be used throughout this appendix.

Comparing the final term of (67) with (11) we wish to swap $(\partial_\mu - \eta_\mu)$ and $(\partial_\nu + \eta_\nu)$ across

sides. This can be done by repeatedly using

$$\begin{aligned}
\partial_\mu f &= f \partial_\mu + [\partial_\mu f] \\
\nabla^2 f &= f \nabla^2 + [\nabla^2 f] + 2[\partial_\alpha f] \partial_\alpha \\
&= f \nabla^2 - [\nabla^2 f] + 2\partial_\alpha [\partial_\alpha f] \\
f \nabla^2 &= \nabla^2 f + [\nabla^2 f] - 2\partial_\alpha [\partial_\alpha f] \\
&= \nabla^2 f - [\nabla^2 f] - 2[\partial_\alpha f] \partial_\alpha
\end{aligned}$$

in the appropriate places.

We first address the terms involving S . These are

$$\begin{aligned}
\partial_\mu S \partial_\nu &= \partial_\nu S \partial_\mu + S_{,\mu} \partial_\nu - \partial_\mu S_{,\nu} + S_{,\mu\nu} \\
\partial_\mu S \eta_\nu &= \eta_\nu S \partial_\mu + [\partial_\mu \eta_\nu S] \\
-\eta_\mu S \partial_\nu &= -\partial_\nu S \eta_\mu + [\partial_\nu \eta_\mu S]
\end{aligned}$$

and thus

$$\hat{R}_{\mu\nu-S}^{1e^-} = (\partial_\mu - \eta_\mu) S (\partial_\nu + \eta_\nu) \quad (D3)$$

$$\begin{aligned}
&= (\partial_\nu + \eta_\nu) S (\partial_\mu - \eta_\mu) \\
&\quad + S_{,\mu} \partial_\nu - \partial_\mu S_{,\nu} + S_{,\mu\nu} \\
&\quad + [\partial_\mu \eta_\nu S] + [\partial_\nu \eta_\mu S].
\end{aligned} \quad (D4)$$

The terms involving the Laplacian are a little more difficult to deal with. Here

$$\begin{aligned}
\partial_\mu \nabla^2 \eta_\nu &= \eta_\nu \nabla^2 \partial_\mu - 2\nabla^2 t_{\mu\nu} \\
&\quad - S_{,\nu} \partial_\mu + 4\eta_\alpha t_{\alpha\nu} \partial_\mu - 4\partial_\alpha t_{\alpha\nu} \partial_\mu \\
-\eta_\mu \nabla^2 \partial_\nu &= -\partial_\nu \nabla^2 \eta_\mu - 2t_{\mu\nu} \nabla^2 \\
&\quad + \partial_\nu S_{,\mu} + 4\partial_\nu t_{\alpha\mu} \eta_\alpha - 4\partial_\nu t_{\alpha\mu} \partial_\alpha
\end{aligned}$$

and

$$\begin{aligned}
\eta_\mu \nabla^2 \eta_\nu &= (\partial_\alpha \eta_\mu - [\partial_\alpha \eta_\mu]) (\eta_\nu \partial_\alpha + [\partial_\alpha \eta_\nu]) \\
&= (\eta_\nu \partial_\alpha + [\partial_\alpha \eta_\nu]) (\partial_\alpha \eta_\mu - [\partial_\alpha \eta_\mu]) \\
&\quad - [\partial_\alpha \eta_\mu] \eta_\nu \partial_\alpha + \partial_\alpha \eta_\mu [\partial_\alpha \eta_\nu] - [\partial_\alpha \eta_\mu] [\partial_\alpha \eta_\nu] \\
&= \eta_\nu \nabla^2 \eta_\mu - 2 [\partial_\alpha \eta_\nu] [\partial_\alpha \eta_\mu] \\
&\quad - \eta_\nu [\nabla^2 \eta_\mu] - [\nabla^2 \eta_\nu] \eta_\mu \\
&\quad - 2 \eta_\nu [\partial_\alpha \eta_\mu] \partial_\alpha + 2 \partial_\alpha [\partial_\alpha \eta_\nu] \eta_\mu \\
&= \eta_\nu \nabla^2 \eta_\mu - 2 [\partial_\alpha \eta_\nu] [\partial_\alpha \eta_\mu] \\
&\quad - \eta_\nu S_{,\mu} - \eta_\mu S_{,\nu} - 4 \eta_\nu t_{\alpha\mu} \eta_\alpha - 4 \eta_\alpha t_{\alpha\nu} \eta_\mu \\
&\quad + 4 \eta_\nu t_{\alpha\mu} \partial_\alpha - 4 \partial_\alpha t_{\alpha\nu} \eta_\mu.
\end{aligned}$$

Combining the above lets us write

$$\begin{aligned}
\hat{R}_{\mu\nu-L}^{1e^-} &= (\partial_\mu - \eta_\mu) \nabla^2 (\partial_\nu + \eta_\nu) \tag{D5} \\
&= (\partial_\nu + \eta_\nu) \nabla^2 (\partial_\mu - \eta_\mu) - 4 \partial_\alpha t_{\mu\nu} \partial_\alpha \\
&\quad - 4 (\partial_\alpha + \eta_\alpha) t_{\alpha\nu} (\partial_\mu - \eta_\mu) \\
&\quad - 4 (\partial_\nu + \eta_\nu) t_{\alpha\mu} (\partial_\alpha - \eta_\alpha) \\
&\quad + 2 [\partial_\alpha \eta_\nu] [\partial_\alpha \eta_\mu] - 2 [\nabla^2 t_{\mu\nu}] \\
&\quad + \partial_\nu S_{,\mu} - S_{,\nu} \partial_\mu + \eta_\nu S_{,\mu} + \eta_\mu S_{,\nu} \tag{D6}
\end{aligned}$$

We can now use (D6) and (D4) together to show

$$\begin{aligned}
4 \hat{R}_{\mu\nu}^{1e^-} &= \hat{R}_{\mu\nu-L}^{1e^-} - \hat{R}_{\mu\nu-S}^{1e^-} \tag{D7} \\
&= (\partial_\nu + \eta_\nu) (\nabla^2 - S) (\partial_\mu - \eta_\mu) \\
&\quad - 4 (\partial_\alpha + \eta_\alpha) t_{\mu\nu} (\partial_\alpha - \eta_\alpha) \\
&\quad - 4 (\partial_\alpha + \eta_\alpha) t_{\alpha\nu} (\partial_\mu - \eta_\mu) \\
&\quad - 4 (\partial_\nu + \eta_\nu) t_{\alpha\mu} (\partial_\alpha - \eta_\alpha) \\
&\quad + S_{,\mu\nu} + 2 [\partial_\alpha \eta_\nu] [\partial_\alpha \eta_\mu] \\
&\quad - 2 [\nabla^2 t_{\mu\nu}] - 4 [\eta_\alpha \partial_\alpha t_{\mu\nu}] \tag{D8}
\end{aligned}$$

where we cancel most terms via

$$\begin{aligned}
\partial_\alpha t_{\mu\nu} \partial_\alpha &= (\partial_\alpha + \eta_\alpha) t_{\mu\nu} (\partial_\alpha - \eta_\alpha) \\
&\quad + [\eta_\alpha \partial_\alpha t_{\mu\nu}] + S t_{\mu\nu} \\
[\partial_\mu \eta_\nu S] &= -2S t_{\mu\nu} + [\eta_\nu S_{,\mu}] \\
[\partial_\nu \eta_\mu S] &= -2S t_{\mu\nu} + [\eta_\mu S_{,\nu}].
\end{aligned}$$

The operator terms are now the same as those of (50). The remaining constant is

$$\begin{aligned}
4K_{\mu\nu} &= S_{,\mu\nu} + 2[\partial_\alpha \eta_\nu][\partial_\alpha \eta_\mu] - 2[\nabla^2 t_{\mu\nu}] - 4[\eta_\alpha \partial_\alpha t_{\mu\nu}] \\
&= S_{,\mu\nu} + 2[\partial_\alpha \eta_\nu][\partial_\alpha \eta_\mu] + S_{,\mu\nu} \\
&\quad - 2[\partial_\nu \eta_\alpha \partial_\mu \eta_\alpha] + 2[\eta_\alpha \partial_\alpha \partial_\mu \eta_\nu] \\
&= 2S_{,\mu\nu} = 4V_{,\mu\nu}
\end{aligned}$$

where we have used $[\nabla^2 t_{\mu\nu}] = -\frac{1}{2}S_{,\mu\nu} + [\partial_\nu \eta_\alpha][\partial_\mu \eta_\alpha] + [\eta_\alpha \partial_\nu \partial_\mu \eta_\alpha]$ and $[\partial_\alpha \eta_\nu] = [\partial_\nu \eta_\alpha]$ to arrive at the final form.

Thus (D8) becomes

$$\begin{aligned}
\hat{R}_{\mu\nu}^{1e^-} &= V_{,\mu\nu} + \frac{1}{4}(\partial_\nu + \eta_\nu)(\nabla^2 - S)(\partial_\mu - \eta_\mu) \\
&\quad - (\partial_\alpha + \eta_\alpha) t_{\mu\nu} (\partial_\alpha - \eta_\alpha) \\
&\quad - (\partial_\alpha + \eta_\alpha) t_{\alpha\nu} (\partial_\mu - \eta_\mu) \\
&\quad - (\partial_\nu + \eta_\nu) t_{\alpha\mu} (\partial_\alpha - \eta_\alpha)
\end{aligned} \tag{D9}$$

which is identical to $\hat{R}_{\mu\nu}^{(\xi)}$ and through it $\hat{R}_{\mu\nu}$ via (45).

Finally, since ψ^1 and through it ϕ^1 and n^1 can be calculated from ξ it is clear that a solution to the CM is identical to a linearised solution of the Schrödinger equation for a one-electron system, and vice versa.

Appendix E: Scalar CM operator in the KS basis

Inserting (50) [and setting $(\nabla^2 - S) = (\partial_\alpha + \eta_\alpha)(\partial_\alpha - \eta_\alpha)$] into (85) and using integration by parts we find

$$\begin{aligned}
R_{jk} = & \int d\mathbf{r} [\hat{D}_\mu \psi_j^*] V_{,\mu\nu} [\hat{D}_\nu \psi_k] \\
& + \frac{1}{4} \int d\mathbf{r} [\hat{D}_\alpha \hat{D}_\mu \hat{D}_\nu \psi_j^*] [\hat{D}_\alpha \hat{D}_\nu \hat{D}_\mu \psi_k] \\
& + \int d\mathbf{r} [\hat{D}_\alpha \hat{D}_\mu \psi_j^*] t_{\mu\nu} [\hat{D}_\alpha \hat{D}_\nu \psi_k] \\
& + \int d\mathbf{r} [\hat{D}_\alpha \hat{D}_\mu \psi_j^*] t_{\alpha\nu} [\hat{D}_\mu \hat{D}_\nu \psi_k] \\
& + \int d\mathbf{r} [\hat{D}_\nu \hat{D}_\mu \psi_j^*] t_{\alpha\mu} [\hat{D}_\alpha \hat{D}_\nu \psi_k]
\end{aligned} \tag{E1}$$

where we have used the shorthand $\hat{D}_\alpha = \partial_\alpha - \eta_\alpha$.

It follows from $\boldsymbol{\eta} = \frac{1}{2} \nabla \log n^0$ that $[\partial_\mu \eta_\nu] = [\partial_\nu \eta_\mu]$ and thus $\partial_\mu \eta_\nu + \eta_\mu \partial_\nu = \eta_\nu \partial_\mu + \partial_\nu \eta_\mu$. Therefore $\hat{D}_\mu \hat{D}_\nu = \hat{D}_\nu \hat{D}_\mu$ and we can simplify (E1) (noting that we can also swap Greek indices as they are summed over) to

$$\begin{aligned}
R_{jk} = & \int d\mathbf{r} [\hat{D}_\mu \psi_j^*] V_{,\mu\nu} [\hat{D}_\nu \psi_k] \\
& + \frac{1}{4} \int d\mathbf{r} [\hat{D}_\alpha \hat{D}_\mu \hat{D}_\nu \psi_j^*] [\hat{D}_\alpha \hat{D}_\mu \hat{D}_\nu \psi_k] \\
& + 3 \int d\mathbf{r} [\hat{D}_\alpha \hat{D}_\mu \psi_j^*] t_{\mu\nu} [\hat{D}_\alpha \hat{D}_\nu \psi_k].
\end{aligned} \tag{E2}$$

REFERENCES

REFERENCES

- ¹E. Madelung, *Naturwissenschaften* **14**, 1004 (1926); *Zeitschrift für Physik A Hadrons and Nuclei* **40**, 322 (1927).
- ²H.-V. Nguyen and S. de Gironcoli, *Phys. Rev. B* **79**, 205114 (2009).
- ³I. V. Tokatly, *Phys. Rev. B* **71**, 165104 (2005); **71**, 165105 (2005).
- ⁴I. V. Tokatly, *Phys. Rev. B* **75**, 125105 (2007).
- ⁵J. Tao, X. Gao, G. Vignale, and I. V. Tokatly, *Phys. Rev. Lett.* **103**, 086401 (2009).
- ⁶X. Gao, J. Tao, G. Vignale, and I. V. Tokatly, *Phys. Rev. B* **81**, 195106 (2010).
- ⁷T. Gould and J. F. Dobson, *Phys. Rev. B* **84**, 241108 (2011).

- ⁸P. Hohenberg and W. Kohn, Phys. Rev. **136**, B864 (1964); W. Kohn and L. J. Sham, **140** (1965).
- ⁹E. Runge and E. K. U. Gross, Phys. Rev. Lett. **52**, 997 (1984).
- ¹⁰See page 1898 of R. H. French, V. A. Parsegian, R. Podgornik, R. F. Rajter, A. Jagota, J. Luo, D. Asthagiri, M. K. Chaudhury, Y.-m. Chiang, S. Granick, S. Kalinin, M. Kardar, R. Kjellander, D. C. Langreth, J. Lewis, S. Lustig, D. Wesolowski, J. S. Wettlaufer, W.-Y. Ching, M. Finnis, F. Houlihan, O. A. von Lilienfeld, C. J. van Oss, and T. Zemb, Rev. Mod. Phys. **82**, 1887 (2010).
- ¹¹J. Harl and G. Kresse, Phys. Rev. B **77**, 045136 (2008); Phys. Rev. Lett. **103**, 056401 (2009).
- ¹²S. Lebègue, J. Harl, T. Gould, J. G. Ángyán, G. Kresse, and J. F. Dobson, Phys. Rev. Lett. **105**, 196401 (2010).
- ¹³H. Eshuis, J. Yarkony, and F. Furche, J. Chem. Phys. **132**, 234114 (2010); H. Eshuis and F. Furche, J. Phys. Chem. Lett. **2**, 983 (2011); H. Eshuis, J. Bates, and F. Furche, Theoretical Chemistry Accounts: Theory, Computation, and Modeling (Theoretica Chimica Acta) **131**, 1 (2012), 10.1007/s00214-011-1084-8.
- ¹⁴J. P. Perdew and K. Schmidt, “Density functional theory and its application to materials,” (Springer, 2001); J. P. Perdew, A. Ruzsinszky, J. Tao, V. N. Staroverov, G. E. Scuseria, and G. I. Csonka, J. Chem. Phys. **123**, 062201 (2005).
- ¹⁵S. Grimme, J. Comp. Chem. **27**, 1787 (2006).
- ¹⁶A. Tkatchenko and M. Scheffler, Phys. Rev. Lett. **102**, 073005 (2009).
- ¹⁷O. A. Vydrov and T. Van Voorhis, Phys. Rev. Lett. **103**, 063004 (2009).
- ¹⁸M. Dion, H. Rydberg, E. Schröder, D. C. Langreth, and B. I. Lundqvist, Phys. Rev. Lett. **92**, 246401 (2004).
- ¹⁹H. Rydberg, B. I. Lundqvist, D. C. Langreth, and M. Dion, Phys. Rev. B **62**, 6997 (2000).
- ²⁰H. Rydberg, M. Dion, N. Jacobson, E. Schröder, P. Hyldgaard, S. I. Simak, D. C. Langreth, and B. I. Lundqvist, Phys. Rev. Lett. **91**, 126402 (2003).
- ²¹D. C. Langreth, M. Dion, H. Rydberg, E. Schröder, P. Hyldgaard, and B. I. Lundqvist, Int. J. Quantum Chem. **101**, 599 (2005).
- ²²J. F. Dobson and T. Gould, Journal of Physics: Condensed Matter **24**, 073201 (2012).
- ²³W. J. Hehre, R. F. Stewart, and J. A. Pople, J. Chem. Phys. **51**, 2657 (1969); M. D. Newton, **51**, 3917 (1969).

- ²⁴J. F. Dobson, Phys. Rev. Lett. **73**, 2244 (1994).
- ²⁵F. Furche, J. Chem. Phys. **129**, 114105 (2008).
- ²⁶T. C. Wallstrom, Phys. Rev. A **49**, 1613 (1994).
- ²⁷For one-electron this is true by construction. In the 1D cases it follows from the fact that in 1D a scalar function can be written as the derivative of another function without the curl.
- ²⁸J. F. Dobson, J. Chem. Phys. **94**, 4328 (1991); **98**, 8870 (1993).
- ²⁹E. Schrödinger, Ann. Phys. (Leipzig) **82**, 265 (1927).
- ³⁰W. Pauli, *Handbuch der Physik*, edited by H. Geiger and K. Scheel (Springer, Berlin, 1933).
- ³¹R. P. Feynman, *Forces and Stresses in Molecules*, B.Sc. Thesis, Massachusetts Institute of Technology (1939).
- ³²P. C. Martin and J. Schwinger, Phys. Rev. **115**, 1342 (1959).
- ³³R. Puff and N. Gillis, Annals of Physics **46**, 364 (1968).
- ³⁴A. Savin and H.-J. Flad, International Journal of Quantum Chemistry **56**, 327 (1995).
- ³⁵T. Leininger, H. Stoll, H.-J. Werner, and A. Savin, Chem. Phys. Lett. **275**, 151 (1997).
- ³⁶I. C. Gerber and J. G. Àngyàn, Chem. Phys. Lett. **415**, 100 (2005).